

Rabbit anti-Human CDK6 Polyclonal Antibody | anti-CDK6 antibody

Phospho-CDK6(Tyr292) Antibody

CATALOG # AAA321423	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CDK6; Polyclonal Antibody; Phospho-CDK6(Tyr292) Antibody; CDK 6; CDK6_HUMAN; Cell division protein kinase 6; Crk 2; Crk2; Cyclin dependent kinase 6; Cyclin-dependent kinase 6; MCPH12; MGC59692; p40; PLSTIRE; Serine/threonine protein kinase PLSTIRE; Serine/threonine-protein kinase PLSTIRE; STQTL11; anti-CDK6 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-CDK6(Tyr292) antibody detects endogenous levels of CDK6.
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	326
Applicable Applications for anti-CDK6 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human CDK6 around the phosphorylation site of Tyr292.
Subcellular Location	Nucleus; Cytoskeleton
Tissue Specificity	Expressed ubiquitously. Accumulates in squamous cell carcinomas, proliferating hematopoietic progenitor cells, beta-cells of pancreatic islets of Langerhans, and neuroblastomas. Reduced levels in differentiating cells.
Predicted Cross Reactivity	Pig, Bovine, Horse, Sheep, Rabbit, Dog, Chicken
Similarity	Pig (100%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (92%), Dog (100%), Chicken (92%)
Conjugation	Unconjugated
Epitope	Phospho Tyr292
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)
AAA321423 at 1/100 staining Human breast cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)
Western blot analysis of extracts of Jurkat cells, using Phospho-CDK6(Tyr292) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CDK6 antibody	Function: Serine/threonine-protein kinase involved in the control of the cell cycle and differentiation; promotes G1/S transition. Phosphorylates pRB/RB1 and NPM1. Interacts with D-type G1 cyclins during interphase at G1 to form a pRB/RB1 kinase and controls the entrance into the cell cycle. Involved in initiation and maintenance of cell cycle exit during cell differentiation; prevents cell proliferation and regulates negatively cell differentiation, but is required for the proliferation of specific cell types (e.g. erythroid and hematopoietic cells). Essential for cell proliferation within the dentate gyrus of the hippocampus and the subventricular zone of the lateral ventricles. Required during thymocyte development. Promotes the production of newborn neurons, probably by modulating G1 length. Promotes, at least in astrocytes, changes in patterns of gene expression, changes in the actin cytoskeleton including loss of stress fibers, and enhanced motility during cell differentiation. Prevents myeloid differentiation by interfering with RUNX1 and reducing its transcription transactivation activity, but promotes proliferation of normal myeloid progenitors. Delays senescence. Promotes the proliferation of beta-cells in pancreatic islets of Langerhans. May play a role in the centrosome organization during the cell cycle phases (PubMed:23918663). Subunit Structure: Interaction with D-type G1 cyclins. Cyclin binding promotes enzyme activation by phosphorylation at Thr-177 (By similarity). Binds to RUNX1, CDKN2D, FBXO7 and CDKN2C/p18-INK4c. Forms a cytoplasmic complex with Hsp90/HSP90AB1 and CDC37. FBXO7-binding promotes D-type cyclin binding. Interacts with Kaposi's sarcoma herpesvirus (KSHV) V-cyclin and herpesvirus saimiri (V-cyclin/ECLF2); the CDK6/V-cyclin complex phosphorylates NPM1 and thus lead to viral reactivation by reducing viral LANA levels. Post-translational Modifications: Thr-177 phosphorylation and Tyr-24 dephosphorylation promotes kinase activity. Similarity: Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	223718134
NCBI GeneID	1021
NCBI Accession #	NP_001138778.1
NCBI GenBank Nucleotide #	NM_001145306.1
UniProt Accession #	Q00534; A4D1G0
Molecular Weight	Observed: 36 kDa Predicted: 37 kDa
NCBI Official Full Name	cyclin-dependent kinase 6
NCBI Official Synonym Full Names	cyclin dependent kinase 6
NCBI Official Symbol	CDK6
NCBI Official Synonym Symbols	MCPH12; PLSTIRE
NCBI Protein Information	cyclin-dependent kinase 6

UniProt Protein Name	Cyclin-dependent kinase 6
UniProt Gene Name	CDK6
UniProt Synonym Gene Names	CDKN6

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cdk6-antibody-immunohistochemistry-western-blot-elisa/321423>