

Rabbit anti-Human, Mouse CD19 Polyclonal Antibody | anti-CD19 antibody

Phospho-CD19 (Tyr531) Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA321533	Rabbit	Human, Mouse	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot

PRODUCT SYNONYMS


CD19; Polyclonal Antibody; Phospho-CD19 (Tyr531) Antibody; B lymphocyte antigen CD19; CD19 molecule; CD19 antigen; Differentiation antigen CD19; Ig kappa chain V-I region HK101; IGKC; Immunoglobulin kappa constant; immunoglobulin kappa light chain; anti-CD19 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-CD19 (Tyr531) antibody detects endogenous levels of CD19 only when phosphorylated at Tyrosine 531
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	557
Applicable Applications for anti-CD19 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide derived from human CD19 around the phosphorylation site of Tyrosine 531
Subcellular Location	Membrane.
Predicted Cross Reactivity	Bovine, Horse, Sheep, Rabbit, Dog
Similarity	Bovine (100%), Horse (100%), Sheep (100%), Rabbit (100%), Dog (100%)
Conjugation	Unconjugated
Epitope	Phospho Tyr531
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



WB (Western Blot)
Western blot analysis of CD19 phosphorylation expression in Serum treated COS7 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



WB (Western Blot)
Western blot analysis of Phospho-CD19 (Tyr531) Antibody expression in Serum treated COS7 cells lysates. The lane on the right is treated with the antigen-specific peptide.



IF (Immunofluorescence)
AAA321533 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CD19 antibody	Description: CD19 a cell surface molecule which assembles with the antigen receptor of B lymphocytes. A critical signal transduction molecule that regulates B lymphocyte development, activation, and differentiation. Function: Assembles with the antigen receptor of B-lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation. Subunit Structure: Forms a complex with CD21, CD81 and CD225 in the membrane of mature B-cells. Interacts with VAV. Interacts with GRB2 and SOS when phosphorylated on Tyr-348 and/or Tyr-378. Interacts with PLCG2 when phosphorylated on Tyr-409. Interacts with LYN. Post-translational Modifications: Phosphorylated on serine and threonine upon DNA damage, probably by ATM or ATR. Phosphorylated on tyrosine following B-cell activation. Phosphorylated on tyrosine residues by LYN.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	296010921
NCBI GeneID	930
NCBI Accession #	NP_001171569.1
NCBI GenBank Nucleotide #	NM_001178098.1
UniProt Accession #	P15391; Q96S68; Q9BRD6; A0N0P9; F5H635

Molecular Weight	Observed: 61 kDa Predicted: 62 kDa
NCBI Official Full Name	B-lymphocyte antigen CD19 isoform 1
NCBI Official Synonym Full Names	CD19 molecule
NCBI Official Symbol	CD19
NCBI Official Synonym Symbols	B4; CVID3
NCBI Protein Information	B-lymphocyte antigen CD19
UniProt Protein Name	B-lymphocyte antigen CD19
UniProt Gene Name	CD19

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd19-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/321533>