

Rabbit Synuclein alpha Polyclonal Antibody | anti-SNCA antibody

Phospho-Synuclein alpha (Tyr136) Antibody


CATALOG # AAA321609	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	--	--

PRODUCT SYNONYMS


Synuclein alpha; Polyclonal Antibody; Phospho-Synuclein alpha (Tyr136) Antibody; Alpha synuclein; Alpha-synuclein; isoform NACP140; alphaSYN; MGC105443; MGC110988; MGC127560; MGC64356; NACP; Non A beta component of AD amyloid; Non A4 component of amyloid; Non A4 component of amyloid precursor; Non-A beta component of AD amyloid; Non-A-beta component of alzheimers disease amyloid; precursor of; Non-A4 component of amyloid precursor; Non-A4 component of amyloid; OTTHUMP00000218549; OTTHUMP00000218551; OTTHUMP00000218552; OTTHUMP00000218553; OTTHUMP00000218554; PARK 1; PARK 4; PARK1; PARK4; Parkinson disease (autosomal dominant; Lewy body) 4; Parkinson disease familial 1; SNCA; Snca synuclein; alpha (non A4 component of amyloid precursor); SYN; Synuclein alpha 140; Synuclein; SYUA_HUMAN; anti-SNCA antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-Synuclein-alpha (Tyr136) antibody detects endogenous levels of Synuclein-alpha only when phosphorylated at Tyrosine 136
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	140
Applicable Applications for anti-SNCA antibody	ELISA, WB (Western Blot)
Immunogen	A synthesized peptide derived from human Synuclein-alpha around the phosphorylation site of Tyrosine 136
Subcellular Location	Cytoplasm, Cytosol. Membrane. Nucleus. Cell Junction, Synapse. Secreted. Membrane-bound in dopaminergic neurons.
Tissue Specificity	Expressed principally in brain but is also expressed in low concentrations in all tissues examined except in liver. Concentrated in presynaptic nerve terminals.
Conjugation	Unconjugated
Epitope	Phospho Tyr136
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



WB (Western Blot)
Western blot analysis of Phospho-Synuclein alpha (Tyr136) expression in various lysates



WB (Western Blot)
Western blot analysis of Synuclein-alpha phosphorylation expression in Mouse brain tissue lysates, The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-SNCA antibody	Description: Alpha-synuclein is a member of the synuclein family, which also includes beta-and gamma-synuclein. Synucleins are abundantly expressed in the brain and alpha-and beta-synuclein inhibit phospholipase D2 selectively. Function: May be involved in the regulation of dopamine release and transport. Induces fibrillization of microtubule-associated protein tau. Reduces neuronal responsiveness to various apoptotic stimuli, leading to a decreased caspase-3 activation. Subunit Structure: Soluble monomer which can form filamentous aggregates. Interacts with UCHL1 (By similarity). Interacts with phospholipase D and histones. Post-translational Modifications: Phosphorylated, predominantly on serine residues. Phosphorylation by CK1 appears to occur on residues distinct from the residue phosphorylated by other kinases. Phosphorylation of Ser-129 is selective and extensive in synucleinopathy lesions. In vitro, phosphorylation at Ser-129 promoted insoluble fibril formation. Phosphorylated on Tyr-125 by a PTK2B-dependent pathway upon osmotic stress.Hallmark lesions of neurodegenerative synucleinopathies contain alpha-synuclein that is modified by nitration of tyrosine residues and possibly by dityrosine cross-linking to generated stable oligomers. Ubiquitinated. The predominant conjugate is the diubiquitinated form (By similarity). Acetylation at Met-1 seems to be important for proper folding and native oligomeric structure. Similarity: The 'non A-beta component of Alzheimer disease amyloid plaque' domain (NAC domain) is involved in fibrils formation. The middle hydrophobic region forms the core of the filaments. The C-terminus may regulate aggregation and determine the diameter of the filaments. Belongs to the synuclein family.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507109
NCBI GeneID	6622
NCBI Accession #	NP_000336.1
NCBI GenBank Nucleotide #	NM_000345.3
UniProt Accession #	P37840; Q13701; Q4JHI3; Q6IAU6; A8K2A4
Molecular Weight	Observed: 18 kDa Predicted: 15 kDa
NCBI Official Full Name	alpha-synuclein isoform NACP140
NCBI Official Synonym Full Names	synuclein alpha
NCBI Official Symbol	SNCA
NCBI Official Synonym Symbols	PD1; NACP; PARK1; PARK4

NCBI Protein Information	alpha-synuclein
UniProt Protein Name	Alpha-synuclein
UniProt Gene Name	SNCA
UniProt Synonym Gene Names	NACP; PARK1; NACP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/synuclein-alpha-antibody-western-blot-elisa/321609>