

Rabbit Paxillin Polyclonal Antibody | anti-PXN antibody

Phospho-Paxillin (Tyr118) Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA321637	Rabbit	Human, Mouse, Rat	ELISA, Western Blot

PRODUCT SYNONYMS


Paxillin; Polyclonal Antibody; Phospho-Paxillin (Tyr118) Antibody; FLJ16691; FLJ23042; Paired box protein Pax 1; PAX 1; PAX1; PAXI_HUMAN; Paxillin alpha; PXN; PXN protein; anti-PXN antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-Paxillin (Tyr118) antibody detects endogenous levels of Paxillin only when phosphorylated at Tyrosine 118
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	591
Applicable Applications for anti-PXN antibody	ELISA, WB (Western Blot)
Immunogen	A synthesized peptide derived from human Paxillin around the phosphorylation site of Tyrosine 118
Subcellular Location	Cytoplasm > Cytoskeleton. Cell Junction > Focal Adhesion.
Predicted Cross Reactivity	Pig, Bovine, Sheep, Rabbit, Dog, Chicken, Xenopus
Similarity	Pig (100%), Bovine (91%), Sheep (91%), Rabbit (100%), Dog (100%), Chicken (100%), Xenopus (100%)
Conjugation	Unconjugated
Epitope	Phospho Tyr118
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



WB (Western Blot)
Western blot analysis of Paxillin phosphorylation expression in EGF treated HeLa whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



WB (Western Blot)
Western blot analysis of Phospho-Paxillin (Tyr118) expression in various lysates

ADDITIONAL INFORMATION

Related Product Information for anti-PXN antibody	Description: PXN is a multi-domain cytoskeletal protein that localizes primarily to focal adhesion sites to the extracellular matrix. Phosphorylated by focal adhesion kinase (FAK) and is a component of integrin signaling. Its phosphorylation provides docking sites for recruitment of signaling molecules to focal adhesions. Function: Cytoskeletal protein involved in actin-membrane attachment at sites of cell adhesion to the extracellular matrix (focal adhesion). Subunit Structure: Binds in vitro to vinculin as well as to the SH3 domain of SRC and, when tyrosine phosphorylated, to the SH2 domain of V-CRK. Isoform beta binds to PTK2/FAK1 but weakly to vinculin. Isoform gamma binds to vinculin but only weakly to PTK2/FAK1. Interacts with GIT1, NUDT16L1/SDOS and TGFBI1. Component of cytoplasmic complexes, which also contain GIT1, ARHGEF6 and PAK1. Interacts with PTK2/FAK1 and PTK2B/PYK2. Binds ASAP2. Interacts with unphosphorylated ITGA4. Interacts with RNF5 and PDCD10. Interacts with NEK3 and this interaction is prolactin-dependent. Interacts with PTK6. Interacts with SORBS1, PARVA and PARVB. Interacts (via cytoplasmic domain) with CEACAM1; the interaction is phosphotyrosyl-dependent (PubMed:11035932). Post-translational Modifications: Phosphorylated by MAPK1/ERK2 (By similarity). Phosphorylated on tyrosine residues during integrin-mediated cell adhesion, embryonic development, fibroblast transformation and following stimulation of cells by mitogens. Phosphorylation at Ser-244 by CDK5 reduces its interaction with PTK2/FAK1 in matrix-cell focal adhesions (MCFA) during oligodendrocytes (OLs) differentiation. Phosphorylation at Tyr-31 and Tyr-118 by PTK6 promote the activation of RAC1 via CRK/CrKII, thereby promoting migration and invasion. Phosphorylation at Ser-250 by SLK is required for PXN redistribution and cell motility (PubMed:23128389). Similarity: Belongs to the paxillin family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	170932514
NCBI GeneID	5829
NCBI Accession #	NP_001074324.1
NCBI GenBank Nucleotide #	NM_001080855.2
UniProt Accession #	P49023; O14970; O14971; O60360; Q5HYA4; B2RAI3; B7ZMB4
Molecular Weight	Observed: 68 kDa Predicted: 65 kDa
NCBI Official Full Name	paxillin isoform 2
NCBI Official Synonym Full Names	paxillin
NCBI Official Symbol	PXN
NCBI Protein Information	paxillin

UniProt Protein Name	Paxillin
UniProt Gene Name	PXN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/paxillin-antibody-western-blot-elisa/321637>