

Rabbit SOX2 Polyclonal Antibody | anti-SOX2 antibody

SOX2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA321821	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SOX2; Polyclonal Antibody; SOX2 Antibody; ANOP3; cb236; Delta EF2a; lcc; MCOPS3; MGC148683; MGC2413; RGD1565646; Sex determining region Y box 2; SOX 2; Sox2; SOX2_HUMAN; SRY (sex determining region Y) box 2; SRY box containing gene 2; SRY related HMG box 2; SRY related HMG box gene 2; SRY-box 2; Transcription factor SOX 2; Transcription factor SOX-2; ysb; anti-SOX2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin (Thermo Fisher Scientific).
Form/Format	Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	~1mg/ml (varies by lot)
Sequence Length	317
Applicable Applications for anti-SOX2 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human SOX2, corresponding to a region within the internal amino acids.
Preparation and Storage	Store at -20°C. Stable for 12 months from date of receipt.

IMAGES




IHC (Immunohistochemistry)

AAA321821 at 1/100 staining Rat spleen tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-Rabbit Ab was used as the secondary Ab.




IHC (Immunohistochemistry)

AAA321821 at 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IF (Immunofluorescence)

AAA321821 staining U-251 MG cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.



WB (Western Blot)

Western blot analysis of SOX2 expression in HUES7 cells. The lane on the left is treated with the antigen-specific peptide.

NCBI AND UNIPROT INFORMATION

NCBI GI #	28195386
NCBI GeneID	6657
NCBI Accession #	NP_003097.1
NCBI GenBank Nucleotide #	NM_003106.3
UniProt Accession #	P48431; Q14537
Molecular Weight	34 kDa
NCBI Official Full Name	transcription factor SOX-2
NCBI Official Synonym Full Names	SRY-box 2
NCBI Official Symbol	SOX2
NCBI Official Synonym Symbols	ANOP3; MCOPS3
NCBI Protein Information	transcription factor SOX-2
UniProt Protein Name	Transcription factor SOX-2
UniProt Gene Name	SOX2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/sox2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/321821>