

Rabbit HSL Polyclonal Antibody | anti-HSL antibody

HSL Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA322385	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HSL; Polyclonal Antibody; HSL Antibody; Hormone sensitive lipase; Hormone sensitive lipase testicular isoform; Hormone-sensitive lipase; LHS; Lipase hormone sensitive; LIPE; LIPS_HUMAN; anti-HSL antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	HSL antibody detects endogenous levels of total HSL
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	1076
Applicable Applications for anti-HSL antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human HSL
Subcellular Location	Cell Membrane. Membrane > Caveola. Cytoplasm > Cytosol. Found in the high-density caveolae. Translocates to the cytoplasm from the caveolae upon insulin stimulation.
Predicted Cross Reactivity	Pig, Bovine, Horse, Sheep, Rabbit, Dog
Similarity	Pig (100%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (91%), Dog (100%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemisry)
AAA322385 at 1/100 staining Human prostate tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)
Western blot analysis of HSL expression in Jurkat whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



IF (Immunofluorescence)
AAA322385 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-HSL antibody	Description: Hormone sensitive lipase is a lipolytic enzyme of the 'GDXG' family. Plays a rate limiting step in triglyceride lipolysis. In adipose tissue and heart, it primarily hydrolyzes stored triglycerides to free fatty acids, while in steroidogenic tissues, it principally converts cholesteryl esters to free cholesterol for steroid hormone production. Function: In adipose tissue and heart, it primarily hydrolyzes stored triglycerides to free fatty acids, while in steroidogenic tissues, it principally converts cholesteryl esters to free cholesterol for steroid hormone production. Subunit Structure: Interacts with CAVIN1 in the adipocyte cytoplasm (PubMed:17026959). Interacts with PLIN5 (By similarity). Post-translational Modifications: Phosphorylation by AMPK may block translocation to lipid droplets. Similarity: Belongs to the 'GDXG' lipolytic enzyme family.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	21328446
NCBI GeneID	3991
NCBI Accession #	NP_005348.2
NCBI GenBank Nucleotide #	NM_005357.3
UniProt Accession #	Q05469; Q3LRT2; Q6NSL7
Molecular Weight	Observed: 85 kDa Predicted: 117 kDa

NCBI Official Full Name	hormone-sensitive lipase
NCBI Official Synonym Full Names	lipase E, hormone sensitive type
NCBI Official Symbol	LIPE
NCBI Official Synonym Symbols	HSL; LHS; AOMS4; FPLD6
NCBI Protein Information	hormone-sensitive lipase
UniProt Protein Name	Hormone-sensitive lipase
UniProt Gene Name	LIPE
UniProt Synonym Gene Names	HSL

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hsl-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/322385>