

Rabbit anti-Human, Mouse ABCG2 Polyclonal Antibody | anti-ABCG2 antibody

ABCG2 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323446	Rabbit	Human, Mouse	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ABCG2; Polyclonal Antibody; ABCG2 Antibody; ABC transporter; ABC15; ABCG 2; ABCG2_HUMAN; ABCP; ATP binding cassette sub family G (WHITE) member 2; ATP binding cassette transporter G2; ATP-binding cassette sub-family G member 2; BCRP; BCRP1; BMDP; Breast cancer resistance protein; CD338; CDw338; CDw338 antigen; EST157481; GOUT1; MGC102821; Mitoxantrone resistance associated protein; Mitoxantrone resistance-associated protein; MRX; Multi drug resistance efflux transport ATP binding cassette sub family G (WHITE) member 2; MXR; MXR1; Placenta specific ATP binding cassette transporter; Placenta specific MDR protein; Placenta-specific ATP-binding cassette transporter; UAQTL1; anti-ABCG2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	ABCG2 antibody detects endogenous levels of total ABCG2
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	611
Applicable Applications for anti-ABCG2 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Cell Membrane
Tissue Specificity	Highly expressed in placenta. Low expression in small intestine, liver and colon.
Conjugation	Unconjugated



WB (Western Blot)

Western blot analysis of ABCG2 expression in ; HepG2 cell lysate, The lane on the left is treated with the antigen-specific peptide.



IHC (Immunohistochemistry)

AAA323446 at 1/100 staining Mouse colon tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-ABCG2 antibody

Description: Xenobiotic transporter that may play an important role in the exclusion of xenobiotics from the brain. May be involved in brain-to-blood efflux. Appears to play a major role in the multidrug resistance phenotype of several cancer cell lines. When overexpressed, the transfected cells become resistant to mitoxantrone, daunorubicin and doxorubicin, display diminished intracellular accumulation of daunorubicin, and manifest an ATP-dependent increase in the efflux of rhodamine 123.

Function: High-capacity urate exporter functioning in both renal and extrarenal urate excretion. Plays a role in porphyrin homeostasis as it is able to mediate the export of protoporphyrin IX (PPIX) both from mitochondria to cytosol and from cytosol to extracellular space, and cellular export of heme, and heme. Xenobiotic transporter that may play an important role in the exclusion of xenobiotics from the brain. Appears to play a major role in the multidrug resistance phenotype of several cancer cell lines. Implicated in the efflux of numerous drugs and xenobiotics: mitoxantrone, the photosensitizer pheophorbide, camptothecin, methotrexate, azidothymidine (AZT), and the anthracyclines daunorubicin and doxorubicin.

Subunit Structure: Monomer under reducing conditions, the minimal functional units is a homodimer; disulfide-linked, but the major oligomeric form in plasma membranes is a homotetramer with possibility of higher order oligomerization up to homododecamers.

Post-translational Modifications: Glycosylation-deficient ABCG2 is normally expressed and functional.

Similarity: The extracellular loop 3 (ECL3) is involved in binding porphyrins and transfer them to other carriers, probably albumin. Belongs to the ABC transporter superfamily. ABCG family. Eye pigment precursor importer (TC 3.A.1.204) subfamily.

NCBI AND UNIPROT INFORMATION

NCBI GI #	383792176
NCBI GeneID	9429
NCBI Accession #	NP_001244315.1
NCBI GenBank Nucleotide #	NM_001257386.2
UniProt Accession #	Q9UNQ0; O95374; Q4W5I3; Q53ZQ1; Q569L4; Q5YLG4; Q86V64; Q8IX16; Q96LD6; Q96TA8; A0A1W3; A8K1T5
Molecular Weight	Observed: 72 kDa Predicted: 73 kDa
NCBI Official Full Name	ATP-binding cassette sub-family G member 2 isoform 2
NCBI Official Synonym Full Names	ATP binding cassette subfamily G member 2 (Junior blood group)
NCBI Official Symbol	ABCG2
NCBI Official Synonym Symbols	MRX; MXR; ABCP; BCRP; BMDP; MXR1; ABC15; BCRP1; CD338; GOUT1; MXR-1; CDw338; UAQTL1; EST157481

NCBI Protein Information	ATP-binding cassette sub-family G member 2
UniProt Protein Name	ATP-binding cassette sub-family G member 2
UniProt Gene Name	ABCG2
UniProt Synonym Gene Names	ABCP; BCRP; BCRP1; MXR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/abcg2-antibody-immunohistochemistry-western-blot-elisa/323446>