

Rabbit anti-Human, Mouse EDG1 Polyclonal Antibody | anti-EDG1 antibody

EDG1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323592	Rabbit	Human, Mouse	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

EDG1; Polyclonal Antibody; EDG1 Antibody; CD363; CHEDG 1; CHEDG1; D1S3362; ECGF 1; ECGF1; EDG 1; endothelial differentiation G protein coupled receptor 1; Endothelial differentiation G-protein coupled receptor 1; Endothelial differentiation sphingolipid G protein coupled receptor 1; FLJ58121; G protein coupled sphingolipid receptor; g protein-coupled receptor edg-1; S1P receptor 1; S1P receptor Edg 1; S1P receptor Edg-1; S1P receptor Edg1; S1P(1) receptor; S1P1; s1pr1; S1PR1_HUMAN; sphingolipid g-protein-coupled receptor 1; Sphingosine 1 phosphate receptor Edg 1; Sphingosine 1 phosphate receptor EDG1; sphingosine 1-phosphate receptor 1; Sphingosine 1-phosphate receptor 1; Sphingosine 1-phosphate receptor Edg-1; Sphingosine 1phosphate receptor type 1 S1P1; anti-EDG1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	EDG1 antibody detects endogenous levels of total EDG1
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	382
Applicable Applications for anti-EDG1 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Cell Membrane
Tissue Specificity	Endothelial cells, and to a lesser extent, in vascular smooth muscle cells, fibroblasts, melanocytes, and cells of epithelioid origin.
Conjugation	Unconjugated

IMAGES



IHC (Immunohistochemistry)

AAA323592 at 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)

Western blot analysis of EDG1 expression in Jurkat cell lysate at 25ug/lane;. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-EDG1 antibody	Description: Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. This inducible epithelial cell G-protein-coupled receptor may be involved in the processes that regulate the differentiation of endothelial cells. Seems to be coupled to the G(i) subclass of heteromeric G proteins. Function: G-protein coupled receptor for the bioactive lysosphingolipid sphingosine 1-phosphate (S1P) that seems to be coupled to the G(i) subclass of heteromeric G proteins. Signaling leads to the activation of RAC1, SRC, PTK2/FAK1 and MAP kinases. Plays an important role in cell migration, probably via its role in the reorganization of the actin cytoskeleton and the formation of lamellipodia in response to stimuli that increase the activity of the sphingosine kinase SPHK1. Required for normal chemotaxis toward sphingosine 1-phosphate. Required for normal embryonic heart development and normal cardiac morphogenesis. Plays an important role in the regulation of sprouting angiogenesis and vascular maturation. Inhibits sprouting angiogenesis to prevent excessive sprouting during blood vessel development. Required for normal egress of mature T-cells from the thymus into the blood stream and into peripheral lymphoid organs. Plays a role in the migration of osteoclast precursor cells, the regulation of bone mineralization and bone homeostasis (By similarity). Plays a role in responses to oxidized 1-palmitoyl-2-arachidonoyl-sn-glycero-3-phosphocholine by pulmonary endothelial cells and in the protection against ventilator-induced lung injury. Subunit Structure: Interacts with GNAI1 and GNAI3. Post-translational Modifications: S1P-induced endothelial cell migration requires the PKB/AKT1-mediated phosphorylation of the third intracellular loop at the Thr-236 residue. Similarity: Belongs to the G-protein coupled receptor 1 family.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	1002741592
NCBI GeneID	1901
NCBI Accession #	NP_001307659.1
NCBI GenBank Nucleotide #	NM_001320730.1
UniProt Accession #	P21453; Q9BYY4; Q9NYN8; D3DT66
Molecular Weight	Observed: 43 kDa Predicted: 43 kDa
NCBI Official Full Name	sphingosine 1-phosphate receptor 1
NCBI Official Synonym Full Names	sphingosine-1-phosphate receptor 1
NCBI Official Symbol	S1PR1
NCBI Official Synonym Symbols	EDG1; S1P1; CD363; ECGF1; EDG-1; CHEDG1; D1S3362
NCBI Protein Information	sphingosine 1-phosphate receptor 1
UniProt Protein Name	Sphingosine 1-phosphate receptor 1
UniProt Gene Name	S1PR1
UniProt Synonym Gene Names	CHEDG1; EDG1; S1P receptor 1; S1P1; S1P receptor Edg-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.