

Rabbit EDNRB Polyclonal Antibody | anti-EDNRB antibody

EDNRB Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323594	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

EDNRB; Polyclonal Antibody; EDNRB Antibody; ABCDS; Ednra; EDNRB_HUMAN; Endothelin B receptor; Endothelin B receptor precursor; Endothelin receptor Non selective type; Endothelin receptor non-selective type; Endothelin receptor type B; ET B; ET-B; ET-BR; ETB; ETBR; ETRB; Hirschsprung disease 2; HSCR; HSCR2; OTTHUMP00000018534; OTTHUMP00000178736; WS4A; anti-EDNRB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	EDNRB antibody detects endogenous levels of total EDNRB
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	442
Applicable Applications for anti-EDNRB antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Cell Membrane
Tissue Specificity	Expressed in placental stem villi vessels, but not in cultured placental villi smooth muscle cells.
Conjugation	Unconjugated

IMAGES



WB (Western Blot)

Western blot analysis of EDNRB expression in HEK293 cells. The lane on the left is treated with the antigen-specific peptide.



IF (Immunofluorescence)

AAA323594 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600 was used as secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-EDNRB antibody	Description: Non-specific receptor for endothelin 1, 2, and 3. Mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. Post-translational Modifications: Palmitoylation of Cys-402 was confirmed by the palmitoylation of Cys-402 in a deletion mutant lacking both Cys-403 and Cys-405. Similarity: Belongs to the G-protein coupled receptor 1 family. Endothelin receptor subfamily. EDNRB sub-subfamily.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557547
NCBI GeneID	1910
NCBI Accession #	NP_000106.1
NCBI GenBank Nucleotide #	NM_000115.4
UniProt Accession #	P24530; O15343; Q59GB1; Q5W0G9; Q8NHM6; Q8NHM7; Q8NHM8; Q8NHM9; Q9UD23; Q9UQK3; A2A2Z8; A8K3T4
Molecular Weight	Observed: 50 kDa Predicted: 50 kDa
NCBI Official Full Name	endothelin receptor type B isoform 1
NCBI Official Synonym Full Names	endothelin receptor type B
NCBI Official Symbol	EDNRB
NCBI Official Synonym Symbols	ETB; ET-B; ETB1; ETBR; ETRB; HSCR; WS4A; ABCDS; ET-BR; HSCR2
NCBI Protein Information	endothelin receptor type B
UniProt Protein Name	Endothelin receptor type B
UniProt Gene Name	EDNRB
UniProt Synonym Gene Names	ET-B; ET-BR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.