

Rabbit CaMKII Polyclonal Antibody | anti-CaMKII antibody

CaMKII Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323655	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CaMKII; Polyclonal Antibody; CaMKII Antibody; Calcium/calmodulin-dependent protein kinase type II subunit alpha; Calcium/calmodulin-dependent protein kinase type II subunit delta; CaM kinase II subunit alpha; CaM kinase II subunit delta; CAM2; CaMK-II subunit alpha; CaMK-II subunit delta; CAMK2A; CAMK2B; CAMKA; CAMKB; CAMKD; CaMKII alpha; KIAA0968; anti-CaMKII antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin (Thermo Fisher Scientific).
Form/Format	Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	~1mg/ml (varies by lot)
Sequence Length	666
Applicable Applications for anti-CaMKII antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human CaMKII, corresponding to a region within the internal amino acids.
Preparation and Storage	Store at -20°C. Stable for 12 months from date of receipt.

IMAGES




IHC (Immunohistochemistry)

AAA323655 at 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)

Western blot analysis of CaMKII using COLO205 whole cell lysates



IF (Immunofluorescence)

AAA323655 staining U87 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

NCBI AND UNIPROT INFORMATION

NCBI GI #	26051204
NCBI GeneID	816
NCBI Accession #	NP_001211.3
NCBI GenBank Nucleotide #	NM_001220.4
UniProt Accession #	Q13554; O95437; O95438; O95599; Q9UGH7; A4D2K0; A4D2K1; A4D2K2; A4D2K3; A4D2K4; A4D2K5; A4D2K6
Molecular Weight	50 kD, 60 kD
NCBI Official Full Name	calcium/calmodulin-dependent protein kinase type II subunit beta isoform 1
NCBI Official Synonym Full Names	calcium/calmodulin dependent protein kinase II beta
NCBI Official Symbol	CAMK2B
NCBI Official Synonym Symbols	CAM2; CAMK2; CAMKB; MRD54
NCBI Protein Information	calcium/calmodulin-dependent protein kinase type II subunit beta
UniProt Protein Name	Calcium/calmodulin-dependent protein kinase type II subunit beta
UniProt Gene Name	CAMK2B
UniProt Synonym Gene Names	CAM2; CAMK2; CAMKB; CaM kinase II subunit beta; CaMK-II subunit beta

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/camkii-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/323655>