

Rabbit Dyskerin Polyclonal Antibody | anti-DKC1 antibody

Dyskerin Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323749	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot

PRODUCT SYNONYMS


Dyskerin; Polyclonal Antibody; Dyskerin Antibody; CBF5; CBF5 homolog; Cbf5p homolog; DKC 1; DKC; Dkc1; DKC1_HUMAN; DKCX; Dyskeratosis congenita 1; Dyskeratosis congenita 1 dyskerin; H/ACA ribonucleoprotein complex subunit 4; NAP 57; NAP57; NOLA 4; NOLA4; Nopp140 associated protein of 57 kDa; Nopp140-associated protein of 57 kDa; Nucleolar protein family A member 4; Nucleolar protein NAP57; snoRNP protein DKC1; XAP 101; XAP101; anti-DKC1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Dyskerin antibody detects endogenous levels of total Dyskerin
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	509
Applicable Applications for anti-DKC1 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide derived from human DKC1, corresponding to a region within the internal amino acids.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



WB (Western Blot)
Western blot analysis of extracts from HeLa cells, using Dyskerin antibody.



IF (Immunofluorescence)
AAA323749 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-DKC1 antibody	Function: Isoform 1: Required for ribosome biogenesis and telomere maintenance. Probable catalytic subunit of H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which catalyzes pseudouridylation of rRNA. This involves the isomerization of uridine such that the ribose is subsequently attached to C5, instead of the normal N1. Each rRNA can contain up to 100 pseudouridine ('psi') residues, which may serve to stabilize the conformation of rRNAs. Also required for correct processing or intranuclear trafficking of TERC, the RNA component of the telomerase reverse transcriptase (TERT) holoenzyme. Subunit Structure: Part of the H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which contains NHP2/NOLA2, GAR1/NOLA1, NOP10/NOLA3, and DKC1/NOLA4, which is presumed to be the catalytic subunit. The complex contains a stable core formed by binding of one or two NOP10-DKC1 heterodimers to NHP2; GAR1 subsequently binds to this core via DKC1. The complex binds a box H/ACA small nucleolar RNA (snoRNA), which may target the specific site of modification within the RNA substrate. During assembly, the complex contains NAF1 instead of GAR1/NOLA1. The complex also interacts with TERC, which contains a 3'-terminal domain related to the box H/ACA snoRNAs. Specific interactions with snoRNAs or TERC are mediated by GAR1 and NHP2. Associates with NOLC1/NOPP140. H/ACA snoRNPs interact with the SMN complex, consisting of SMN1 or SMN2, GEMIN2/SIP1, DDX20/GEMIN3, and GEMIN4. This is mediated by interaction between GAR1 and SMN1 or SMN2. The SMN complex may be required for correct assembly of the H/ACA snoRNP complex. Component of the telomerase holoenzyme complex at least composed of TERT, DKC1, WRAP53/TCAB1, NOP10, NHP2, GAR1, TEP1, EST1A, POT1 and a telomerase RNA template component (TERC). Interacts with SHQ1; this interaction may lead to the stabilization of DKC1, from the time of its synthesis until its association with NOP10, NHP2, and NAF1 at the nascent H/ACA RNA. Similarity: Belongs to the pseudouridine synthase TruB family.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	215599015
NCBI GeneID	1736
NCBI Accession #	NP_001135935.1
NCBI GenBank Nucleotide #	NM_001142463.2
UniProt Accession #	O60832; O43845; Q96G67; Q9Y505; F5BSB3
Molecular Weight	Observed: 57 kD Predicted: 58 kDa
NCBI Official Full Name	H/ACA ribonucleoprotein complex subunit DKC1 isoform 2
NCBI Official Synonym Full Names	dyskerin pseudouridine synthase 1
NCBI Official Symbol	DKC1

NCBI Official Synonym Symbols	DKC; CBF5; DKCX; NAP57; NOLA4; XAP101
NCBI Protein Information	H/ACA ribonucleoprotein complex subunit DKC1
UniProt Protein Name	H/ACA ribonucleoprotein complex subunit DKC1
UniProt Gene Name	DKC1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/dyskerin-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/323749>