

Rabbit hnRNP M Polyclonal Antibody | anti-HNRNPM antibody

hnRNP M Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323904	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot



PRODUCT SYNONYMS

hnRNP M; Polyclonal Antibody; hnRNP M Antibody; CEA receptor; CEAR; Heterogeneous nuclear ribonucleoprotein M; Heterogenous nuclear ribonucleoprotein M4; hnRNA binding protein M4; Hnrnrm; HNRNPM4; HNRPM; HNRPM_HUMAN; HNRPM4; HTGR1; M3; M4; M4 protein; N-acetylglucosamine receptor 1; NAGR1; anti-HNRNPM antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	hnRNP M antibody detects endogenous levels of total hnRNP M
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	730
Applicable Applications for anti-HNRNPM antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Nucleus, Nucleolus.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)
AAA323904 at 1/100 staining Human breast cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

WB (Western Blot)
Western blot analysis of extracts from HT-29 cells, using hnRNP M antibody. The lane on the left is treated with the antigen-specific peptide.



IF (Immunofluorescence)
AAA323904 staining HT29 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-HNRNPM antibody	Function: Pre-mRNA binding protein in vivo, binds avidly to poly(G) and poly(U) RNA homopolymers in vitro. Involved in splicing. Acts as a receptor for carcinoembryonic antigen in Kupffer cells, may initiate a series of signaling events leading to tyrosine phosphorylation of proteins and induction of IL-1 alpha, IL-6, IL-10 and tumor necrosis factor alpha cytokines. Subunit Structure: Identified in the spliceosome C complex. Post-translational Modifications: Sumoylated.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	14141152
NCBI GeneID	4670
NCBI Accession #	NP_005959.2
NCBI GenBank Nucleotide #	NM_005968.4
UniProt Accession #	P52272; Q15584; Q8WZ44; Q96H56; Q9BWL9; Q9Y492
Molecular Weight	Observed: 85; 78 kD Predicted: 78 kDa
NCBI Official Full Name	heterogeneous nuclear ribonucleoprotein M isoform a
NCBI Official Synonym Full Names	heterogeneous nuclear ribonucleoprotein M

NCBI Official Symbol	HNRNPM
NCBI Official Synonym Symbols	CEAR; HNRPM; HTGR1; NAGR1; HNRPM4; HNRNPM4; hnRNP M
NCBI Protein Information	heterogeneous nuclear ribonucleoprotein M
UniProt Protein Name	Heterogeneous nuclear ribonucleoprotein M
UniProt Gene Name	HNRNPM
UniProt Synonym Gene Names	HNRPM; NAGR1; hnRNP M

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hnrm-p-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/323904>