

Rabbit TFAM Polyclonal Antibody | anti-TFAM antibody

TFAM Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323934	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry



PRODUCT SYNONYMS

TFAM; Polyclonal Antibody; TFAM Antibody; anscription factor 6-like 1; Mitochondrial transcription factor 1; mitochondrial transcription factor A; MtTF1; mtTFA; TCF 6; TCF-6; TCF6; TCF6L1; TCF6L2; TCF6L3; TFAM_HUMAN; Transcription factor 6; Transcription factor 6 like 2 (mitochondrial transcription factor); Transcription factor 6 like 2; Transcription factor 6-like 2; transcription factor 6-like 3; Transcription factor A; mitochondrial; mitochondrial precursor; anti-TFAM antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	TFAM antibody detects endogenous levels of total TFAM
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	214
Applicable Applications for anti-TFAM antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry)
Immunogen	A synthesized peptide
Subcellular Location	Mitochondrion.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)
AAA323934 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

IHC (Immunohistochemistry)
AAA323934 at 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-TFAM antibody	Function: Binds to the mitochondrial light strand promoter and functions in mitochondrial transcription regulation (PubMed:29445193). Required for accurate and efficient promoter recognition by the mitochondrial RNA polymerase. Promotes transcription initiation from the HSP1 and the light strand promoter by binding immediately upstream of transcriptional start sites. Is able to unwind DNA. Bends the mitochondrial light strand promoter DNA into a U-turn shape via its HMG boxes. Required for maintenance of normal levels of mitochondrial DNA. May play a role in organizing and compacting mitochondrial DNA. Subunit Structure: Monomer; binds DNA as a monomer. Upon metabolic stress, forms a complex composed of FOXO3, SIRT3, TFAM and POLRMT (PubMed:29445193). Interacts with TFB1M and TFB2M. Interacts with CLPX; this enhances DNA-binding. Post-translational Modifications: Phosphorylation by PKA within the HMG box 1 impairs DNA binding and promotes degradation by the AAA+ Lon protease. Similarity: Binds DNA via its HMG boxes. When bound to the mitochondrial light strand promoter, bends DNA into a U-turn shape, each HMG box bending the DNA by 90 degrees.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	399498566
NCBI GeneID	7019
NCBI Accession #	NP_001257711.1
NCBI GenBank Nucleotide #	NM_001270782.1
UniProt Accession #	Q00059; Q5U0C6; A8MRB2; A9QXC6; B5BU05
Molecular Weight	Predicted: 30 kDa
NCBI Official Full Name	transcription factor A, mitochondrial isoform 2
NCBI Official Synonym Full Names	transcription factor A, mitochondrial
NCBI Official Symbol	TFAM
NCBI Official Synonym Symbols	TCF6; MTTF1; MTTFA; TCF6L1; TCF6L2; TCF6L3; MTDPS15
NCBI Protein Information	transcription factor A, mitochondrial
UniProt Protein Name	Transcription factor A, mitochondrial
UniProt Gene Name	TFAM
UniProt Synonym Gene Names	TCF6; TCF6L2; mtTFA; MtTF1; TCF-6

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tfam-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-elisa/323934>