

Rabbit p97 MAPK Polyclonal Antibody | anti-MAPK6 antibody

p97 MAPK Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA324000	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

p97 MAPK; Polyclonal Antibody; p97 MAPK Antibody; ERK-3; ERK3; Extracellular signal regulated kinase 3; Extracellular signal regulated kinase p97; Extracellular signal-regulated kinase 3; HsT17250; MAP kinase 6; MAP kinase isoform p97; MAPK 6; MAPK6; Mitogen activated protein kinase 6; Mitogen-activated protein kinase 6; MK06_HUMAN; p97-MAPK; PRKM6; Protein kinase mitogen activated 5; Protein kinase mitogen activated 6; anti-MAPK6 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	p97 MAPK antibody detects endogenous levels of total p97 MAPK
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	721
Applicable Applications for anti-MAPK6 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Nucleus
Tissue Specificity	Highest expression in the skeletal muscle, followed by the brain. Also found in heart, placenta, lung, liver, pancreas, kidney and skin fibroblasts.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)
AAA324000 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



WB (Western Blot)
Western blot analysis of extracts from K562 cells, using p97 MAPK antibody. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK6 antibody	Function: Atypical MAPK protein. Phosphorylates microtubule-associated protein 2 (MAP2) and MAPKAPK5. The precise role of the complex formed with MAPKAPK5 is still unclear, but the complex follows a complex set of phosphorylation events: upon interaction with atypical MAPKAPK5, ERK3/MAPK6 is phosphorylated at Ser-189 and then mediates phosphorylation and activation of MAPKAPK5, which in turn phosphorylates ERK3/MAPK6. May promote entry in the cell cycle (By similarity). Subunit Structure: Heterodimer with ERK4/MAPK4. Interacts with (via FRIEDE motif) MAPKAPK5 (By similarity). Interacts with UBE3A; this interaction may be indirect and mediated by HERC2, possibly via HERC2 interaction with NEURL4. Post-translational Modifications: Phosphorylated at Ser-189 by PAK1, PAK2 and PAK3 resulting in catalytic activation. Phosphorylated by MAPKAPK5 at other sites. Ubiquitination at Met-1 leads to degradation by the proteasome pathway. Similarity: In contrast to classical MAPKs, the TXY motif within the activation loop is replaced by the SEG motif, whose phosphorylation activates the MAP kinases. Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506091
NCBI GeneID	5597
NCBI Accession #	NP_002739.1
NCBI GenBank Nucleotide #	NM_002748.3
UniProt Accession #	Q16659; Q68DH4; Q8IYN8; B2R945; B5BU65
Molecular Weight	Observed: 85 kD Predicted: 83 kDa
NCBI Official Full Name	mitogen-activated protein kinase 6
NCBI Official Synonym Full Names	mitogen-activated protein kinase 6
NCBI Official Symbol	MAPK6
NCBI Official Synonym Symbols	ERK3; PRKM6; p97MAPK; HsT17250
NCBI Protein Information	mitogen-activated protein kinase 6
UniProt Protein Name	Mitogen-activated protein kinase 6
UniProt Gene Name	MAPK6

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/p97-mapk-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/324000>