

Rabbit anti-Human, Mouse MKNK1 Polyclonal Antibody | anti-MKNK1 antibody

MKNK1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA324111	Rabbit	Human, Mouse	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

MKNK1; Polyclonal Antibody; MKNK1 Antibody; MAP kinase interacting kinase 1; MAP kinase interacting serine/threonine kinase 1; MAP kinase signal integrating kinase 1; MAP kinase signal-integrating kinase 1; MAP kinase-interacting serine/threonine-protein kinase 1; MAPK signal integrating kinase 1; MITOGEN-ACTIVATED PROTEIN KINASE-INTERACTING SERINE/THREONINE KINASE 1; mknk1; MKNK1_HUMAN; MNK 1; Mnk1; anti-MKNK1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	MKNK1 antibody detects endogenous levels of total MKNK1
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	424
Applicable Applications for anti-MKNK1 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Cytoplasm and Cytoplasm. Nucleus.
Tissue Specificity	Ubiquitous.
Predicted Cross Reactivity	Pig, Zebrafish, Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus
Similarity	Pig (100%), Zebrafish (100%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (100%), Dog (100%), Chicken (100%), Xenopus (100%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)

AAA324111 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.



WB (Western Blot)

Western blot analysis of extracts from HuvEc cells, using MKNK1 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-MKNK1 antibody	Function: May play a role in the response to environmental stress and cytokines. Appears to regulate translation by phosphorylating EIF4E, thus increasing the affinity of this protein for the 7-methylguanosine-containing mRNA cap. Subunit Structure: Interacts with the C-terminal regions of EIF4G1 and EIF4G2. Also binds to dephosphorylated ERK1 and ERK2, and to the p38 kinases. Post-translational Modifications: Dual phosphorylation of Thr-250 and Thr-255 activates the kinase. Phosphorylation of Thr-385 activates the kinase. MAPK3/ERK1 is one of the kinases which activate MKNK1/MNK1. Phosphorylation by PAK2 leads to a reduced phosphorylation of EIF4G1. Similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	207113186
NCBI GeneID	8569
NCBI Accession #	NP_001129025.1
NCBI GenBank Nucleotide #	NM_001135553.2
UniProt Accession #	Q9BUB5; O00312; Q5TC06; Q5TC07; Q6V0N6; D3DQ20; D3DQ21
Molecular Weight	Observed: 51 kD Predicted: 52 kDa
NCBI Official Full Name	MAP kinase-interacting serine/threonine-protein kinase 1 isoform 3
NCBI Official Synonym Full Names	MAP kinase interacting serine/threonine kinase 1
NCBI Official Symbol	MKNK1
NCBI Official Synonym Symbols	MNK1
NCBI Protein Information	MAP kinase-interacting serine/threonine-protein kinase 1
UniProt Protein Name	MAP kinase-interacting serine/threonine-protein kinase 1
UniProt Gene Name	MKNK1
UniProt Synonym Gene Names	MNK1; MAPK signal-integrating kinase 1; Mnk1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

