

Rabbit PDHA1 Polyclonal Antibody | anti-PDHA1 antibody

PDHA1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA325021	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

PDHA1; Polyclonal Antibody; PDHA1 Antibody; ODP_A_HUMAN; PDH; PDHA; PDHCE1A; PDHE1 A type I; PDHE1-A type I; PHE1A; Pyruvate Dehydrogenase (lipoamide) alpha 1; Pyruvate dehydrogenase complex; E1 alpha polypeptide 1; Pyruvate Dehydrogenase E1 alpha; Pyruvate dehydrogenase E1 component subunit alpha; somatic form; mitochondrial; anti-PDHA1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	PDHA1 antibody detects endogenous levels of total PDHA1
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	390
Applicable Applications for anti-PDHA1 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human PDHA1, corresponding to a region within C-terminal amino acids.
Conjugation	Unconjugated
Preparation and Storage	Store at -20°C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)

Western Blot analysis of extracts from various samples, using PDHA1 Antibody. Lane 1: Sp2/0 treated with blocking peptide. Lane 2: Sp2/0; Lane 3: 293; Lane 4: HeLa



IF (Immunofluorescence)

AAA325021 staining A549 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



IHC (Immunohistochemistry)

AAA325021 at 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-PDHA1 antibody	The pyruvate dehydrogenase complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO ₂ , and thereby links the glycolytic pathway to the tricarboxylic cycle.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505685
NCBI GeneID	5160
NCBI Accession #	NP_000275.1
NCBI GenBank Nucleotide #	NM_000284.3
UniProt Accession #	P08559; Q53H41; Q5JPT8; Q9NP12; Q9UBJ8; Q9UBU0; Q9UNG4; Q9UNG5; A5YVE9; B2R5P7; B7Z3T7; B7Z3X5
Molecular Weight	43 KD
NCBI Official Full Name	pyruvate dehydrogenase E1 component subunit alpha, somatic form, mitochondrial isoform 1
NCBI Official Synonym Full Names	pyruvate dehydrogenase E1 alpha 1 subunit
NCBI Official Symbol	PDHA1
NCBI Official Synonym Symbols	PDHA; PDHAD; PHE1A; PDHCE1A
NCBI Protein Information	pyruvate dehydrogenase E1 component subunit alpha, somatic form, mitochondrial
UniProt Protein Name	Pyruvate dehydrogenase E1 component subunit alpha, somatic form, mitochondrial
UniProt Gene Name	PDHA1
UniProt Synonym Gene Names	PHE1A

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pdha1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/325021>