

Rabbit anti-Human Cytochrome P450 2D6 Polyclonal Antibody | anti-CYP2D6 antibody

Cytochrome P450 2D6 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA325387	Rabbit	Human	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

Cytochrome P450 2D6; Polyclonal Antibody; Cytochrome P450 2D6 Antibody; CPD6; CYP2D; CYP2D6; CYP2D7AP; CYP2D7BP; CYP2D7P2; CYP2D8P2; CYP2DL1; CYP11D6; Cytochrome P450 DB1; Cytochrome P450 family 2 subfamily D polypeptide 6; Debrisoquine 4 hydroxylase; Flavoprotein linked monooxygenase; Microsomal monooxygenase; P450 DB1; P450C2D; P450DB1; Xenobiotic monooxygenase; anti-CYP2D6 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Cytochrome P450 2D6 antibody detects endogenous levels of total Cytochrome P450 2D6
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	497
Applicable Applications for anti-CYP2D6 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Endoplasmic Reticulum Membrane; Peripheral Membrane Protein. Microsome Membrane; Peripheral Membrane Protein.
Tissue Specificity	By pregnancy.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)
AAA325387 staining HT29 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.



WB (Western Blot)
Western blot analysis of extracts from 293/Jurkat/K562 cells, using CYP2D6 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CYP2D6 antibody	Function: Responsible for the metabolism of many drugs and environmental chemicals that it oxidizes. It is involved in the metabolism of drugs such as antiarrhythmics, adrenoceptor antagonists, and tricyclic antidepressants. Similarity: Belongs to the cytochrome P450 family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	392513721
NCBI GeneID	1565
NCBI Accession #	NP_000097.3
NCBI GenBank Nucleotide #	NM_000106.6
UniProt Accession #	P10635; Q16752; Q2XND6; Q2XND7; Q2XNE0; Q6B012; Q6NXU8
Molecular Weight	Observed: 56 kD Predicted: 56 kDa
NCBI Official Full Name	cytochrome P450 2D6 isoform 1
NCBI Official Synonym Full Names	cytochrome P450 family 2 subfamily D member 6
NCBI Official Symbol	CYP2D6
NCBI Official Synonym Symbols	CPD6; CYP2D; CYP2DL1; CYPIID6; P450C2D; P450DB1; CYP2D7AP; CYP2D7BP; CYP2D7P2; CYP2D8P2; P450-DB1
NCBI Protein Information	cytochrome P450 2D6
UniProt Protein Name	Cytochrome P450 2D6
UniProt Gene Name	CYP2D6
UniProt Synonym Gene Names	CYP2DL1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.