

Rabbit anti-Human GPR18 Polyclonal Antibody | anti-GPR18 antibody

GPR18 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA325520	Rabbit	Human	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

GPR18; Polyclonal Antibody; GPR18 Antibody; G protein coupled receptor 18; GPCRW; GPR 18; N arachidonyl glycine receptor; NAGly receptor; anti-GPR18 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	GPR18 antibody detects endogenous levels of total GPR18
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	331
Applicable Applications for anti-GPR18 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Cell Membrane; Multipass Membrane Protein.
Tissue Specificity	Expressed in midpiece of spermatozoon (at protein level) (PubMed:27572937). Most abundant in testis and spleen (PubMed:16844083). Highly expressed in CD4 and CD8-positive T-cells as well as CD19-positive B-cells (PubMed:16844083).
Predicted Cross Reactivity	Horse, Rabbit, Dog
Similarity	Horse (82%), Rabbit (82%), Dog (80%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)
AAA325520 staining HT29 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



WB (Western Blot)
Western blot analysis of extracts from HT-29 cells, using GPR18 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-GPR18 antibody	Function: Receptor for endocannabinoid N-arachidonyl glycine (NAGly) (PubMed:16844083, PubMed:24762058, PubMed:27572937). However, conflicting results about the role of NAGly as an agonist are reported (PubMed:27018161). Can also be activated by plant-derived and synthetic cannabinoid agonists (PubMed:24762058). The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase (PubMed:16844083). May contribute to regulation of the immune system. Is required for normal homeostasis of CD8+ subsets of intraepithelial lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs)in small intestine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Plays a role in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:27572937). Similarity: Belongs to the G-protein coupled receptor 1 family.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	148235349
NCBI GeneID	2841
NCBI Accession #	NP_001091670.1
NCBI GenBank Nucleotide #	NM_001098200.1
UniProt Accession #	Q14330; Q6GTM3; Q96HI6; Q9H2L2
Molecular Weight	Observed: 35 kD Predicted: 39 kDa
NCBI Official Full Name	N-arachidonyl glycine receptor
NCBI Official Synonym Full Names	G protein-coupled receptor 18
NCBI Official Symbol	GPR18
NCBI Protein Information	N-arachidonyl glycine receptor
UniProt Protein Name	N-arachidonyl glycine receptor
UniProt Gene Name	GPR18
UniProt Synonym Gene Names	GPCRW; NAGly receptor

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/gpr18-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/325520>