

Rabbit IFN-gamma Polyclonal Antibody | anti-IFNG antibody

IFN-gamma Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA325848	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

IFN-gamma; Polyclonal Antibody; IFN-gamma Antibody; IFG; IFI; IFN gamma; IFN; immune; IFNG; IFNG_HUMAN; Immune interferon; Interferon gamma; anti-IFNG antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Interferon gamma Antibody detects endogenous levels of total Interferon gamma.
Purity/Purification	Peptide affinity purification
Form/Format	IgG in phosphate buffered saline , pH 7.4, 150mMNaCl, 0.02% sodium azide and 50% glycerol
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-IFNG antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human Interferon gamma, corresponding to a region within the internal amino acids.
Preparation and Storage	Store at -20°C.

IMAGES



IHC (Immunohistochemistry)

AAA325848 at 1/100 staining Human mammary cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



WB (Western Blot)

Western blot analysis of mouse brain lysate using IFN-gamma antibody. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-IFNG antibody	Interferons (IFNs) appear both locally and systematically early after viral infection and participate in limiting the spread of infection. They also affect cell differentiation, growth, surface antigen expression and immunoregulation (1). There are three naturally occurring interferons: α , β and γ . IFN- γ is derived from lymphoblastic tissue and has a number of therapeutic applications in the treatment of various human cancers and diseases of viral origin. Recombinant IFN- γ from both natural and synthetic genes binds to a common cell surface receptor and induces antiviral activity in a variety of cell lines. When binding to discrete cell surface receptors on target cells, IFN- γ induces rapid changes in Jak/Stat phosphorylation, which initiates the Jak/Stat signaling pathway
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	56786138
NCBI GeneID	3458
NCBI Accession #	NP_000610.2
NCBI GenBank Nucleotide #	NM_000619.2
UniProt Accession #	P01579; Q53ZV4; B5BU88
Molecular Weight	19kDa
NCBI Official Full Name	interferon gamma
NCBI Official Synonym Full Names	interferon gamma
NCBI Official Symbol	IFNG
NCBI Official Synonym Symbols	IFG; IFI
NCBI Protein Information	interferon gamma
UniProt Protein Name	Interferon gamma
UniProt Gene Name	IFNG
UniProt Synonym Gene Names	IFN-gamma

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ifn-gamma-antibody-immunohistochemistry-western-blot-elisa/325848>