

Rabbit KRT20 Polyclonal Antibody | anti-KRT20 antibody

KRT20 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA325853	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot



PRODUCT SYNONYMS

KRT20; Polyclonal Antibody; KRT20 Antibody; CD20; CK 20; CK-20; CK20; Cytokeratin-20; Cytokeratin20; K1C20_HUMAN; K20; KA20; Keratin 20; keratin 20; type I; keratin 21; rat; homolog of; Keratin; Keratin type I cytoskeletal 20; Keratin-20; Keratin20; KRT 20; KRT 21; KRT21; MGC35423; OTTHUMP00000164518; Protein IT; type I cytoskeletal 20; anti-KRT20 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	KRT20 antibody detects endogenous levels of total KRT20
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	424
Applicable Applications for anti-KRT20 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human KRT20
Subcellular Location	Cytoplasm.
Tissue Specificity	Expressed predominantly in the intestinal epithelium. Expressed in luminal cells of colonic mucosa. Also expressed in the Merkel cells of keratinized oral mucosa; specifically at the tips of some rete ridges of the gingival mucosa, in the basal layer of the palatal mucosa and in the taste buds of lingual mucosa.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES




IF (Immunofluorescence)

AAA325853 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.

IHC (Immunohistochemistry)

AAA325853 at 1/100 staining Mouse colon tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)

Western blot analysis of extracts from Hela using KRT20 Antibody. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT20 antibody

Description: Intermediate-sized filament (IF) protein designated cytokeratin (CK) 20 is a major cellular protein of mature enterocytes and goblet cells commonly found in mucosal epithelium of the mammalian gastrointestinal tract (1). Results strongly suggest that transcriptional regulation of keratin genes in the intestinal epithelium occurs at the level of both immature and terminally differentiated epithelial cells, and is tightly regulated during both fetal development and crypt-to-villus differentiation of the intestinal epithelium (2). CK20 has recently been reported to be useful to distinguish between primary and metastatic lung adenocarcinoma. CK20 expression was significantly more prevalent in adenocarcinoma that originated in the GI tract than that of pulmonary or breast origin (3). Function: Plays a significant role in maintaining keratin filament organization in intestinal epithelia. When phosphorylated, plays a role in the secretion of mucin in the small intestine (By similarity). Subunit Structure: Heterotetramer of two type I and two type II keratins. Associates with KRT8. Post-translational Modifications: Hyperphosphorylation at Ser-13 occurs during the early stages of apoptosis but becomes less prominent during the later stages. Phosphorylation at Ser-13 also increases in response to stress brought on by cell injury (By similarity). Proteolytically cleaved by caspases during apoptosis. Cleavage occurs at Asp-228. Similarity: Belongs to the intermediate filament family.

NCBI AND UNIPROT INFORMATION

NCBI GI #	27894337
NCBI GeneID	54474
NCBI Accession #	NP_061883.1

NCBI GenBank Nucleotide #	NM_019010.2
UniProt Accession #	P35900 ; B2R6W7
Molecular Weight	Observed: 48 kDa Predicted: 49 kDa
NCBI Official Full Name	keratin, type I cytoskeletal 20
NCBI Official Synonym Full Names	keratin 20
NCBI Official Symbol	KRT20
NCBI Official Synonym Symbols	K20; CD20; CK20; CK-20; KRT21
NCBI Protein Information	keratin, type I cytoskeletal 20
UniProt Protein Name	Keratin, type I cytoskeletal 20
UniProt Gene Name	KRT20
UniProt Synonym Gene Names	CK-20; K20

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/krt20-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/325853>