

Rabbit PSMB1 Polyclonal Antibody | anti-PSMB1 antibody

PSMB1 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA325974	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

PSMB1; Polyclonal Antibody; PSMB1 Antibody; FLJ25321; HC5; KIAA1838; Macropain subunit C5; Multicatalytic endopeptidase complex subunit C5; Proteasome (prosome macropain) subunit beta type 1; Proteasome beta 1 subunit; Proteasome component C5; Proteasome gamma chain; Proteasome subunit beta type 1; Proteasome subunit beta type-1; Proteasome subunit HC5; PSB1_HUMAN; PSC5; PSMB 1; anti-PSMB1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	PSMB1 antibody detects endogenous levels of total PSMB1
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	241
Applicable Applications for anti-PSMB1 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human PSMB1
Subcellular Location	Cytoplasm. Nucleus.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



WB (Western Blot)

Western blot analysis of A431 lysate using PRKCD antibody. The lane on the left is treated with the antigen-specific peptide.



IHC (Immunohistochemistry)

AAA325974 at 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMB1 antibody

Description: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is tightly linked to the TBP (TATA-binding protein) gene in human and in mouse, and is transcribed in the opposite orientation in both species. [provided by RefSeq, Jul 2008]

Function: Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).

Subunit Structure: The 26S proteasome consists of a 20S proteasome core and two 19S regulatory subunits. The 20S proteasome core is a barrel-shaped complex made of 28 subunits that are arranged in four stacked rings. The two outer rings are each formed by seven alpha subunits, and the two inner rings are formed by seven beta subunits. The proteolytic activity is exerted by three beta-subunits PSMB5, PSMB6 and PSMB7. Interacts with SERPINB2 (PubMed:14732874). Interacts with HIV-1 TAT protein (PubMed:14550573).

Similarity: Belongs to the peptidase T1B family.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506193
NCBI GeneID	5689
NCBI Accession #	NP_002784.1
NCBI GenBank Nucleotide #	NM_002793.3
UniProt Accession #	P20618; Q9BWA8; B5BU76
Molecular Weight	Observed: 26 kDa Predicted: 27 kDa
NCBI Official Full Name	proteasome subunit beta type-1
NCBI Official Synonym Full Names	proteasome subunit beta 1

NCBI Official Symbol	PSMB1
NCBI Official Synonym Symbols	HC5; PSC5; PMSB1
NCBI Protein Information	proteasome subunit beta type-1
UniProt Protein Name	Proteasome subunit beta type-1
UniProt Gene Name	PSMB1
UniProt Synonym Gene Names	PSC5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/psmb1-antibody-immunohistochemistry-western-blot-elisa/325974>