

Rabbit DUSP6 Polyclonal Antibody | anti-DUSP6 antibody

DUSP6 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326063	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

DUSP6; Polyclonal Antibody; DUSP6 Antibody; Dual specificity phosphatase 6; Dual specificity phosphatase 6 isoform a; Dual specificity protein phosphatase 6; Dual specificity protein phosphatase PYST1; DUS6_HUMAN; DUSP 6; DUSP 6a; Dusp6; DUSP6a; HH19; MAP kinase phosphatase 3; Mitogen activated protein kinase phosphatase 3; Mitogen-activated protein kinase phosphatase 3; MKP 3; MKP-3; MKP3; PYST 1; PYST1; Serine/threonine specific protein phosphatase; anti-DUSP6 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	DUSP6 antibody detects endogenous levels of total DUSP6
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	381
Applicable Applications for anti-DUSP6 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human DUSP6
Subcellular Location	Cytoplasm.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.


IMAGES



IHC (Immunohistochemisry)
AAA326063 at 1/100 staining Human brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IF (Immunofluorescence)
AAA326063 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600 was used as secondary antibody.



WB (Western Blot)
Western blot analysis of 3T3 lysate usingDUSP6 antibody. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-DUSP6 antibody	Description: MAP kinases are inactivated by dual-specificity protein phosphatases (DUSP) that differ in their substrate specificity, tissue distribution, inducibility by extracellular stimuli and cellular localization. DUSPs, also known as MAPK phosphatases (MKP), specifically dephosphorylate both threonine and tyrosine residues in MAPK P-loops and have been shown to play important roles in regulating the function of the MAPK family (1, 2). At least 13 members of the family (DUSP1-10, DUSp14, DUSP16, and DUSP22) display unique substrate specificities for various MAP kinases (3). MAPK phosphatases typically contain an amino-terminal rhodanese-fold responsible for DUSP docking to MAPK family members and a carboxy-terminal catalytic domain (4). These phosphatases can play important roles in development, immune system function, stress responses and metabolic homeostasis (5), and also in the development of cancer and the response of cancer cells to chemotherapy (6).DUSP6 specifically dephosphorylates ERK MAP kinase (7). Function: Inactivates MAP kinases. Has a specificity for the ERK family (PubMed:9858808). Plays an important role in alleviating chronic postoperative pain. Necessary for the normal dephosphorylation of the long-lasting phosphorylated forms of spinal MAPK1/3 and MAP kinase p38 induced by peripheral surgery, which drives the resolution of acute postoperative allodynia (By similarity). Also important for dephosphorylation of MAPK1/3 in local wound tissue, which further contributes to resolution of acute pain (By similarity). Similarity: Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	42764683
NCBI GeneID	1848

NCBI Accession #	NP_001937.2
NCBI GenBank Nucleotide #	NM_001946.3
UniProt Accession #	Q16828 ; O75109 ; Q53Y75 ; Q9BSH6
Molecular Weight	Observed: 42 kDa Predicted: 43 kDa
NCBI Official Full Name	dual specificity protein phosphatase 6 isoform a
NCBI Official Synonym Full Names	dual specificity phosphatase 6
NCBI Official Symbol	DUSP6
NCBI Official Synonym Symbols	HH19; MKP3; PYST1
NCBI Protein Information	dual specificity protein phosphatase 6
UniProt Protein Name	Dual specificity protein phosphatase 6
UniProt Gene Name	DUSP6
UniProt Synonym Gene Names	MKP3; PYST1; MAP kinase phosphatase 3; MKP-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/dusp6-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/326063>