

Rabbit ATF3 Polyclonal Antibody | anti-ATF3 antibody

ATF3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326425	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ATF3; Polyclonal Antibody; ATF3 Antibody; Activating transcription factor 3; ATF3_HUMAN; ATF3deltaZip2; ATF3deltaZip2c; ATF3deltaZip3; cAMP dependent transcription factor ATF3; cAMP-dependent transcription factor ATF-3; Cyclic AMP dependent transcription factor ATF3; Cyclic AMP-dependent transcription factor ATF-3; anti-ATF3 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	ATF3 antibody detects endogenous levels of total ATF3
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	181
Applicable Applications for anti-ATF3 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human ATF3
Subcellular Location	Nucleus.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)

AAA326425 at 1/100 staining Mouse liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)

Western blot analysis of extracts of various tissue, using ATF3 antibody.



IF (Immunofluorescence)

AAA326425 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-ATF3 antibody	Description: This gene encodes a member of the mammalian activation transcription factor/cAMP responsive element-binding (CREB) protein family of transcription factors. This gene is induced by a variety of signals, including many of those encountered by cancer cells, and is involved in the complex process of cellular stress response. Multiple transcript variants encoding different isoforms have been found for this gene. It is possible that alternative splicing of this gene may be physiologically important in the regulation of target genes. Function: This protein binds the cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Represses transcription from promoters with ATF sites. It may repress transcription by stabilizing the binding of inhibitory cofactors at the promoter. Isoform 2 activates transcription presumably by sequestering inhibitory cofactors away from the promoters. Subunit Structure: Binds DNA as a homodimer or a heterodimer. Similarity: Belongs to the bZIP family. ATF subfamily.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	71902536
NCBI GeneID	467
NCBI Accession #	NP_001025458.1
NCBI GenBank Nucleotide #	NM_001030287.3
UniProt Accession #	P18847; Q5VTZ2; Q6ICQ9; Q7Z566; Q8WYM6
Molecular Weight	Observed: 20 kDa Predicted: 21 kDa
NCBI Official Full Name	cyclic AMP-dependent transcription factor ATF-3 isoform 1
NCBI Official Synonym Full Names	activating transcription factor 3
NCBI Official Symbol	ATF3
NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-3
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-3
UniProt Gene Name	ATF3
UniProt Synonym Gene Names	cAMP-dependent transcription factor ATF-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.