

Rabbit WDR5 Polyclonal Antibody | anti-WDR5 antibody

WDR5 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326522	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

WDR5; Polyclonal Antibody; WDR5 Antibody; 2410008O07Rik; AA408785; AA960360; BIG 3; Big; BIG3; BMP2 induced 3 kb gene protein; BMP2 induced gene; 3-KB; BMP2-induced 3-kb gene protein; MGC114572; OTTHUMP00000162494; SWD 3; SWD3; SWD3 Set1c WD40 repeat protein homolog; WD repeat containing protein 5; WD repeat domain 5; WD repeat protein 5; WD repeat-containing protein 5; WD repeat-containing protein BIG-3; WDR 5; Wdr5; WDR5_HUMAN; anti-WDR5 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	WDR5 antibody detects endogenous levels of total WDR5
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	334
Applicable Applications for anti-WDR5 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human WDR5
Subcellular Location	Nucleus.
Predicted Cross Reactivity	Pig, Bovine, Horse, Dog
Similarity	Pig (83%), Bovine (100%), Horse (100%), Dog (100%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)
AAA326522 at 1/100 staining Rat brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)
Western blot analysis of Hela whole cell lysates, using WDR5 Antibody. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-WDR5 antibody

Description: This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-aspartic acid (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This protein contains 7 WD repeats. Alternatively spliced transcript variants encoding the same protein have been identified.

Function: Contributes to histone modification. May position the N-terminus of histone H3 for efficient trimethylation at 'Lys-4'. As part of the MLL1/MLL complex it is involved in methylation and dimethylation at 'Lys-4' of histone H3. H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation. As part of the NSL complex it may be involved in acetylation of nucleosomal histone H4 on several lysine residues. May regulate osteoblasts differentiation.

Subunit Structure: Interacts with PAXBP1; the interaction is direct and links a WDR5-containing histone methyltransferase complex to PAX7 and PAX3. Interacts with HCFC1. Component of the ATAC complex, a complex with histone acetyltransferase activity on histones H3 and H4. Component of the SET1 complex, at least composed of the catalytic subunit (SETD1A or SETD1B), WDR5, WDR82, RBBP5, ASH2L/ASH2, CXXC1/CFP1, HCFC1 and DPY30. Core component of several methyltransferase-containing complexes including MLL1/MLL, MLL2/3 (also named ASCOM complex) and MLL4/WBP7. Each complex is at least composed of ASH2L, RBBP5, WDR5, DPY30, one or more specific histone methyltransferases (KMT2A/MLL1, KMT2D/MLL2, KMT2C/MLL3 and KMT2B/MLL4), and the facultative components PAGR1, BAP18, CHD8, E2F6, HCFC1, HCFC2, HSP70, INO80C, KDM6A, KANSL1, LAS1L, MAX, MCRC1, MEN1, MGA, MYST1/MOF, NCOA6, PAXIP1/PTIP, PELP1, PHF20, PRP31, RING2, RUVB1/TIP49A, RUVB2/TIP49B, SENP3, TAF1, TAF4, TAF6, TAF7, TAF9, TEX10 and alpha-and beta-tubulin. Component of the NSL complex at least composed of MOF/KAT8, KANSL1, KANSL2, KANSL3, MCRC1, PHF20, OGT1/OGT, WDR5 and HCFC1. Interacts with KMT2A/MLL1 and RBBP5; the interaction is direct. Component of the ADA2A-containing complex (ATAC), composed of KAT14, KAT2A, TADA2L, TADA3L, ZZ3, MBIP, WDR5, YEATS2, CCDC101 and DR1. In the complex, it probably interacts directly with KAT2A, MBIP and KAT14. Interacts with histone H3. Interacts with SETD1A. Part of a complex composed at least of ASCL2, EMSY, HCFC1, HSPA8, CCAR2, MATR3, MKI67, RBBP5, TUBB2A, WDR5 and ZNF335; this complex may have a histone H3-specific methyltransferase activity. Interacts with PER1.

Similarity: Belongs to the WD repeat WDR5/wds family.

NCBI AND UNIPROT INFORMATION

NCBI GI #	16554627
NCBI GeneID	11091
NCBI Accession #	NP_060058.1
NCBI GenBank Nucleotide #	NM_017588.2
UniProt Accession #	P61964; Q91VA5; Q9NWX7; Q9UGP9

Molecular Weight	Observed: 36 kDa Predicted: 37 kDa
NCBI Official Full Name	WD repeat-containing protein 5
NCBI Official Synonym Full Names	WD repeat domain 5
NCBI Official Symbol	WDR5
NCBI Official Synonym Symbols	SWD3; BIG-3; CFAP89
NCBI Protein Information	WD repeat-containing protein 5
UniProt Protein Name	WD repeat-containing protein 5
UniProt Gene Name	WDR5
UniProt Synonym Gene Names	BIG3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/wdr5-antibody-immunohistochemistry-western-blot-elisa/326522>