

Rabbit ICAM2 Polyclonal Antibody | anti-ICAM2 antibody

ICAM2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326534	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ICAM2; Polyclonal Antibody; ICAM2 Antibody; CD102; CD102 antigen; ICAM 2; Intercellular adhesion molecule 2; Ly60; anti-ICAM2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	ICAM2 antibody detects endogenous levels of total ICAM2
Purity/Purification	Peptide affinity purification
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	275
Applicable Applications for anti-ICAM2 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human ICAM2, corresponding to a region within the internal amino acids.
Subcellular Location	Membrane; Single-Pass Type I Membrane Protein
Conjugation	Unconjugated
Important	For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)

AAA326534 at 1/100 staining Human urothelial cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)

Western blot analysis of extracts of Jurkat, using ICAM2 antibody. The lane on the left is treated with the antigen-specific peptide.



IF (Immunofluorescence)

AAA326534 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-ICAM2 antibody	Description: The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008] Function: ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. Subunit Structure: Interacts with RDX, EZR and MSN. Similarity: Belongs to the immunoglobulin superfamily. ICAM family.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	153082696
NCBI GeneID	3384
NCBI Accession #	NP_000864.2
NCBI GenBank Nucleotide #	NM_000873.3
UniProt Accession #	P13598; Q14600
Molecular Weight	31 kDa
NCBI Official Full Name	intercellular adhesion molecule 2
NCBI Official Synonym Full Names	intercellular adhesion molecule 2
NCBI Official Symbol	ICAM2
NCBI Official Synonym Symbols	CD102
NCBI Protein Information	intercellular adhesion molecule 2
UniProt Protein Name	Intercellular adhesion molecule 2
UniProt Gene Name	ICAM2
UniProt Synonym Gene Names	ICAM-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.