

Rabbit TNFRSF11B Polyclonal Antibody | anti-TNFRSF11B antibody

TNFRSF11B Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326584	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

TNFRSF11B; Polyclonal Antibody; TNFRSF11B Antibody; MGC29565; OCIF; OPG; Osteoclastogenesis inhibitory factor; Osteoprotegerin; PDB5; TNF receptor superfamily member 11b; TNFRSF 11B; TR 1; TR1; TR11B_HUMAN; Tumor necrosis factor receptor superfamily member 11B; anti-TNFRSF11B antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	TNFRSF11B antibody detects endogenous levels of total TNFRSF11B
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	401
Applicable Applications for anti-TNFRSF11B antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human TNFRSF11B
Subcellular Location	Secreted.
Tissue Specificity	Highly expressed in adult lung, heart, kidney, liver, spleen, thymus, prostate, ovary, small intestine, thyroid, lymph node, trachea, adrenal gland, testis, and bone marrow. Detected at very low levels in brain, placenta and skeletal muscle. Highly expressed in fetal kidney, liver and lung.
Predicted Cross Reactivity	Pig, Bovine, Horse, Sheep, Rabbit, Dog, Chicken
Similarity	Pig (100%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (100%), Dog (100%), Chicken (80%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)

Western blot analysis of K562 whole cell lysates, using TNFRSF11B Antibody. The lane on the left is treated with the antigen-specific peptide.



IHC (Immunohistochemistry)

AAA326584 at 1/100 staining Mouse testis tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IF (Immunofluorescence)

AAA326584 staining 293 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-TNFRSF11B antibody

Description: The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein is an osteoblast-secreted decoy receptor that functions as a negative regulator of bone resorption. This protein specifically binds to its ligand, osteoprotegerin ligand, both of which are key extracellular regulators of osteoclast development. Studies of the mouse counterpart also suggest that this protein and its ligand play a role in lymph-node organogenesis and vascular calcification. Alternatively spliced transcript variants of this gene have been reported, but their full length nature has not been determined. [provided by RefSeq, Jul 2008]
Function: Acts as decoy receptor for TNFSF11/RANKL and thereby neutralizes its function in osteoclastogenesis. Inhibits the activation of osteoclasts and promotes osteoclast apoptosis in vitro. Bone homeostasis seems to depend on the local ratio between TNFSF11 and TNFRSF11B. May also play a role in preventing arterial calcification. May act as decoy receptor for TNFSF10/TRAIL and protect against apoptosis. TNFSF10/TRAIL binding blocks the inhibition of osteoclastogenesis.
Subunit Structure: Homodimer. Interacts with TNFSF10 and TNFSF11.
Post-translational Modifications: N-glycosylated. Contains sialic acid residues. The N-terminus is blocked.

NCBI AND UNIPROT INFORMATION

NCBI GI #	148743793
NCBI GeneID	4982
NCBI Accession #	NP_002537.3
NCBI GenBank Nucleotide #	NM_002546.3
UniProt Accession #	O00300; O60236; Q53FX6; Q9UHP4; B2R9A8
Molecular Weight	Observed: 46 kDa Predicted: 47 kDa
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 11B
NCBI Official Synonym Full Names	TNF receptor superfamily member 11b
NCBI Official Symbol	TNFRSF11B
NCBI Official Synonym Symbols	OPG; TR1; OCIF; PDB5
NCBI Protein Information	tumor necrosis factor receptor superfamily member 11B
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 11B

UniProt Gene Name	TNFRSF11B
UniProt Synonym Gene Names	OCIF; OPG

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tnfrsf11b-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/326584>