

Rabbit NDUFS1 Polyclonal Antibody | anti-NDUFS1 antibody

NDUFS1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326790	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

NDUFS1; Polyclonal Antibody; NDUFS1 Antibody; CI-75kD; Complex I 75kD; Complex I; mitochondrial respiratory chain; 75 kD subunit; Complex I-75kD; mitochondrial; NADH coenzyme Q reductase; NADH dehydrogenase (ubiquinone) FeS protein 1 (75kD) (NADH coenzyme Q reductase); NADH ubiquinone oxidoreductase 75 kDa subunit mitochondrial; NADH-ubiquinone oxidoreductase 75 kDa subunit; NDUS1_HUMAN; PRO1304; anti-NDUFS1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	NDUFS1 antibody detects endogenous levels of total NDUFS1
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	691
Applicable Applications for anti-NDUFS1 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human NDUFS1
Subcellular Location	Mitochondrion Inner Membrane.
Predicted Cross Reactivity	Pig, Zebrafish, Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus
Similarity	Pig (100%), Zebrafish (100%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (100%), Dog (100%), Chicken (100%), Xenopus (100%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)

Western blot analysis of HeLa whole cell lysates, using NDUFS1 Antibody. The lane on the left is treated with the antigen-specific peptide.



IF (Immunofluorescence)

AAA326790 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



IHC (Immunohistochemistry)

AAA326790 at 1/100 staining Human gastric tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-NDUFS1 antibody

Description: The protein encoded by this gene belongs to the complex I 75 kDa subunit family. Mammalian complex I is composed of 45 different subunits. It locates at the mitochondrial inner membrane. This protein has NADH dehydrogenase activity and oxidoreductase activity. It transfers electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone. This protein is the largest subunit of complex I and it is a component of the iron-sulfur (IP) fragment of the enzyme. It may form part of the active site crevice where NADH is oxidized. Mutations in this gene are associated with complex I deficiency. Several transcript variants encoding different isoforms have been found for this gene.

Function: Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity). This is the largest subunit of complex I and it is a component of the iron-sulfur (IP) fragment of the enzyme. It may form part of the active site crevice where NADH is oxidized.

Subunit Structure: Complex I is composed of 45 different subunits.

Similarity: Belongs to the complex I 75 kDa subunit family.

NCBI AND UNIPROT INFORMATION

NCBI GI #	316983154
NCBI GeneID	4719
NCBI Accession #	NP_001186910.1
NCBI GenBank Nucleotide #	NM_001199981.1
UniProt Accession #	P28331; Q53TR8; Q8N1C4; Q8TCC9; B4DIN9; B4DJA0; B4DPG1; B4DUC1; E7ENF3
Molecular Weight	Observed: 79 kDa Predicted: 80 kDa
NCBI Official Full Name	NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial isoform 2
NCBI Official Synonym Full Names	NADH:ubiquinone oxidoreductase core subunit S1
NCBI Official Symbol	NDUFS1
NCBI Official Synonym Symbols	CI-75k; CI-75Kd; PRO1304
NCBI Protein Information	NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial
UniProt Protein Name	NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial
UniProt Gene Name	NDUFS1
UniProt Synonym Gene Names	CI-75kD

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.