

Rabbit CXCR2 Polyclonal Antibody | anti-CXCR2 antibody

CXCR2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326836	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

CXCR2; Polyclonal Antibody; CXCR2 Antibody; C-X-C chemokine receptor type 2; CD 182; CD182; CD182 antigen; CDw128b; Chemokine (CXC) receptor 2; CMKAR2; CXC-R2; CXCR 2; CXCR-2; CXCR2_HUMAN; GRO/MGSA receptor; High affinity interleukin-8 receptor B; IL 8 receptor type 2; IL 8R B; IL-8 receptor type 2; IL-8R B; IL8 RB; IL8 receptor type 2; IL8R B; IL8R2; IL8RA; Interleukin 8 Receptor B; Interleukin 8 receptor; beta; type 2; anti-CXCR2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	CXCR2 antibody detects endogenous levels of total CXCR2
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	360
Applicable Applications for anti-CXCR2 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthetic peptide of human CXCR2
Subcellular Location	Cell Membrane
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)
AAA326836 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



WB (Western Blot)
Western blot analysis of extracts of HT-29, using CXCR2antibody. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-CXCR2 antibody	Description: The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. Function: Receptor for interleukin-8 which is a powerful neutrophil chemotactic factor. Binding of IL-8 to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. Binds to IL-8 with high affinity. Also binds with high affinity to CXCL3, GRO/MGSA and NAP-2. Post-translational Modifications: Phosphorylated upon ligand binding; which is required for desensitization. Similarity: Belongs to the G-protein coupled receptor 1 family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	269973859
NCBI GeneID	3579
NCBI Accession #	NP_001161770.1
NCBI GenBank Nucleotide #	NM_001168298.1
UniProt Accession #	P25025; Q8IUZ1; Q9P2T6; Q9P2T7
Molecular Weight	Observed: 41 kDa Predicted: 41 kDa
NCBI Official Full Name	C-X-C chemokine receptor type 2
NCBI Official Synonym Full Names	C-X-C motif chemokine receptor 2
NCBI Official Symbol	CXCR2
NCBI Official Synonym Symbols	CD182; IL8R2; IL8RA; IL8RB; CMKAR2; CDw128b
NCBI Protein Information	C-X-C chemokine receptor type 2
UniProt Protein Name	C-X-C chemokine receptor type 2
UniProt Gene Name	CXCR2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cxcr2-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/326836>