

Rabbit anti-Human, Mouse MAG Polyclonal Antibody | anti-MAG antibody

MAG Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA327276	Rabbit	Human, Mouse	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MAG; Polyclonal Antibody; MAG Antibody; GMA; MAG_HUMAN; Myelin associated glycoprotein; Myelin-associated glycoprotein; S MAG; S-MAG; Sialic acid binding Ig like lectin 4A; Sialic acid binding immunoglobulin like lectin 4A; Siglec 4a; Siglec-4a; SIGLEC4A; SPG75; anti-MAG antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	MAG antibody detects endogenous levels of total MAG
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	601
Applicable Applications for anti-MAG antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human MAG
Subcellular Location	Membrane.
Tissue Specificity	Both isoform 1 and isoform 2 are detected in myelinated structures in the central and peripheral nervous system, in periaxonal myelin and at Schmidt-Lanterman incisures (PubMed:9495552, PubMed:6200494). Detected in optic nerve, in oligodendroglia and in periaxonal myelin sheaths (PubMed:6200494). Detected in compact myelin (at protein level) (PubMed:6200494). Both isoform 1 and isoform 2 are detected in the central and peripheral nervous system (PubMed:9495552).
Predicted Cross Reactivity	Pig, Bovine, Horse, Sheep, Rabbit, Dog
Similarity	Pig (100%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (100%), Dog (92%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)
Western blot analysis of MAG using COLO205 whole lysates.



IHC (Immunohistochemistry)
AAA327276 at 1/100 staining Human liver cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-MAG antibody	Function: Adhesion molecule that mediates interactions between myelinating cells and neurons by binding to neuronal sialic acid-containing gangliosides and to the glycoproteins RTN4R and RTN4RL2 (By similarity). Not required for initial myelination, but seems to play a role in the maintenance of normal axon myelination. Protects motoneurons against apoptosis, also after injury; protection against apoptosis is probably mediated via interaction with neuronal RTN4R and RTN4RL2. Required to prevent degeneration of myelinated axons in adults; this probably depends on binding to gangliosides on the axon cell membrane (By similarity). Negative regulator of neurite outgrowth; in dorsal root ganglion neurons the inhibition is mediated primarily via binding to neuronal RTN4R or RTN4RL2 and to a lesser degree via binding to neuronal gangliosides. In cerebellar granule cells the inhibition is mediated primarily via binding to neuronal gangliosides. In sensory neurons, inhibition of neurite extension depends only partially on RTN4R, RTN4RL2 and gangliosides. Inhibits axon longitudinal growth (By similarity). Inhibits axon outgrowth by binding to RTN4R (By similarity). Preferentially binds to alpha-2, 3-linked sialic acid. Binds ganglioside Gt1b (By similarity). Subunit Structure: Monomer and homodimer (By similarity). Interacts (via the first three N-terminal Ig-like domains) with RTN4R and RTN4RL2 (By similarity). Interacts with RTN4R (PubMed:19052207). Post-translational Modifications: N-glycosylated. Phosphorylated on tyrosine residues. Similarity: Belongs to the immunoglobulin superfamily. SIGLEC (sialic acid binding Ig-like lectin) family.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	312836851
NCBI GeneID	4099
NCBI Accession #	NP_001186145.1
NCBI GenBank Nucleotide #	NM_001199216.1
UniProt Accession #	P20916; Q15489; Q567S4; B7Z2E5; F5GYC0
Molecular Weight	Observed: 63 kDa Predicted: 70 kDa
NCBI Official Full Name	myelin-associated glycoprotein isoform c
NCBI Official Synonym Full Names	myelin associated glycoprotein
NCBI Official Symbol	MAG
NCBI Official Synonym Symbols	GMA; S-MAG; SPG75; SIGLEC4A; SIGLEC-4A
NCBI Protein Information	myelin-associated glycoprotein
UniProt Protein Name	Myelin-associated glycoprotein

UniProt Gene Name	MAG
UniProt Synonym Gene Names	GMA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mag-antibody-immunohistochemistry-western-blot-elisa/327276>