

Rabbit F5 Polyclonal Antibody | anti-F5 antibody

F5 Antibody

CATALOG # AAA327440	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

F5; Polyclonal Antibody; F5 Antibody; Activated protein C cofactor; APC cofactor; coagulation factor V (proaccelerin; labile factor); Coagulation factor V; coagulation factor V jinjiang A2 domain; Coagulation factor V light chain; FA5_HUMAN; Factor V Leiden; FactorV; FVL; Labile factor; PCCF; Proaccelerin; proaccelerin; labile factor; Protein C cofactor; RPRGL1; THPH2; anti-F5 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin.
Form/Format	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	~1mg/ml (varies by lot)
Applicable Applications for anti-F5 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human Factor V, corresponding to a region within C-terminal amino acids.
Important	For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.
Preparation and Storage	Store at -20 °C. Stable for 12 months from date of receipt.

IMAGES




IHC (Immunohistochemistry)

AAA327440 at 1/100 staining Human melanoma tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



IHC (Immunohistochemistry)

AAA327440 at 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-Rabbit Ab was used as the secondary Ab.



WB (Western Blot)
Western blot analysis F5 using A549 whole cell lysates

NCBI AND UNIPROT INFORMATION

NCBI GI #	105990535
NCBI Accession #	NP_000121.2
NCBI GenBank Nucleotide #	NM_000130.4
UniProt Accession #	P12259; Q14285; Q2EHR5; Q5R346; Q5R347; Q6UPU6; Q8WWQ6; A8K6E8
Molecular Weight	250 kDa
NCBI Official Full Name	coagulation factor V preproprotein
NCBI Official Synonym Full Names	coagulation factor V
NCBI Official Symbol	F5
NCBI Official Synonym Symbols	FVL; PCCF; THPH2; RPRGL1
NCBI Protein Information	coagulation factor V
UniProt Protein Name	Coagulation factor V
UniProt Gene Name	F5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/f5-antibody-immunohistochemistry-western-blot-elisa/327440>