

Rabbit anti-Human, Mouse ADAMTS1 Polyclonal Antibody | anti-ADAMTS1 antibody

ADAMTS1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA327824	Rabbit	Human, Mouse	ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ADAMTS1; Polyclonal Antibody; ADAMTS1 Antibody; A disintegrin and metalloproteinase with thrombospondin motifs 1; ADAM-TS 1; ADAM-TS1; ADAMTS 1; ADAMTS; ADAMTS-1; Adamts1; ATS1_HUMAN; C3 C5; KIAA1346; ; METH 1; METH-1; anti-ADAMTS1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	ADAMTS1 antibody detects endogenous levels of total ADAMTS1
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	967
Applicable Applications for anti-ADAMTS1 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human ADAMTS1
Subcellular Location	Secreted > Extracellular Space > Extracellular Matrix.
Predicted Cross Reactivity	Pig, Zebrafish, Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus
Similarity	Pig (100%), Zebrafish (88%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (100%), Dog (100%), Chicken (88%), Xenopus (100%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)

Western blot analysis of ADAMTS1 using K562 whole cell lysates



IF (Immunofluorescence)

AAA327824 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.



IHC (Immunohistochemistry)

AAA327824 at 1/100 staining Mouse kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-ADAMTS1 antibody	Function: Cleaves aggrecan, a cartilage proteoglycan, at the '1938-Glu-I-Leu-1939' site (within the chondroitin sulfate attachment domain), and may be involved in its turnover (By similarity). Has angiogenic inhibitor activity. Active metalloprotease, which may be associated with various inflammatory processes as well as development of cancer cachexia. May play a critical role in follicular rupture. Post-translational Modifications: The precursor is cleaved by a furin endopeptidase. Glycosylated. Can be O-fucosylated by POFUT2 on a serine or a threonine residue found within the consensus sequence C1-X2-(S/T)-C2-G of the TSP type-1 repeat domains where C1 and C2 are the first and second cysteine residue of the repeat, respectively. Fucosylated repeats can then be further glycosylated by the addition of a beta-1, 3-glucose residue by the glucosyltransferase, B3GALT1. Fucosylation mediates the efficient secretion of ADAMTS family members. Also can be C-glycosylated with one or two mannose molecules on tryptophan residues within the consensus sequence W-X-X-W of the TPRs, and N-glycosylated. These other glycosylations can also facilitate secretion (By similarity). Similarity: The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix. The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	50845384
NCBI GeneID	9510
NCBI Accession #	NP_008919.3
NCBI GenBank Nucleotide #	NM_006988.4
UniProt Accession #	Q9UHI8; Q9NSJ8; Q9P2K0; Q9UH83; Q9UP80; D3DSD5
Molecular Weight	Observed: 105 kDa Predicted: 106 kDa
NCBI Official Full Name	A disintegrin and metalloproteinase with thrombospondin motifs 1 preproprotein
NCBI Official Synonym Full Names	ADAM metalloproteinase with thrombospondin type 1 motif 1
NCBI Official Symbol	ADAMTS1
NCBI Official Synonym Symbols	C3-C5; METH1
NCBI Protein Information	A disintegrin and metalloproteinase with thrombospondin motifs 1
UniProt Protein Name	A disintegrin and metalloproteinase with thrombospondin motifs 1
UniProt Gene Name	ADAMTS1
UniProt Synonym Gene Names	KIAA1346; METH1; ADAM-TS 1; ADAM-TS1; ADAMTS-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.