

Rabbit IRE1 Polyclonal Antibody | anti-IRE1 antibody

IRE1 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA328581	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Western Blot

PRODUCT SYNONYMS

IRE1; Polyclonal Antibody; IRE1 Antibody; Endoplasmic reticulum (ER) to nucleus signalling 1; Endoplasmic reticulum to nucleus signaling 1; Endoplasmic reticulum-to-nucleus signaling 1; Endoribonuclease; ER to nucleus signaling 1; ERN 1; Ern1; ERN1_HUMAN; hIRE 1p; hIRE1p; Inositol requiring 1; S. cerevisiae; homolog of; Inositol requiring enzyme 1; Inositol requiring protein 1; inositol-requiring enzyme 1; Inositol-requiring protein 1; IRE 1; IRE 1a; IRE 1P; Ire1 alpha; Ire1-alpha; IRE1a; Ire1alpha; IRE1P; MGC163277; MGC163279; Protein kinase/endoribonuclease; RGD1559716; Serine/threonine protein kinase/endoribonuclease IRE1; anti-IRE1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid; Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	~1mg/ml (varies by lot)
Sequence Length	977
Applicable Applications for anti-IRE1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide derived from human IRE1, corresponding to a region within the internal amino acids.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



IF (Immunofluorescence)

AAA328581 staining HeLa cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab (AAA328581 1:200) and mouse anti-beta tubulin Ab (1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG (H+L) Ab (Red) and an AlexaFluor488 conjugated goat anti-mouse IgG (H+L) Ab (Green) were used as the secondary antibody. The nuclear counter stain is DAPI (blue).



WB (Western Blot)

Western blot analysis of IRE1 in lysates of K562, using IRE1 Antibody.

NCBI AND UNIPROT INFORMATION

NCBI GI #	193806335
NCBI GeneID	2081
UniProt Accession #	O75460; Q76MJ5
Molecular Weight	110 kDa
NCBI Official Full Name	Serine/threonine-protein kinase/endoribonuclease IRE1
NCBI Official Synonym Full Names	endoplasmic reticulum to nucleus signaling 1
NCBI Official Symbol	ERN1
NCBI Official Synonym Symbols	IRE1; IRE1P; IRE1a; hIRE1p
NCBI Protein Information	serine/threonine-protein kinase/endoribonuclease IRE1
UniProt Protein Name	Serine/threonine-protein kinase/endoribonuclease IRE1
UniProt Gene Name	ERN1
UniProt Synonym Gene Names	hIRE1p; IRE1a
UniProt Entry Name	ERN1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.