

Rabbit Frizzled 4 Polyclonal Antibody | anti-FZD4 antibody

Frizzled 4 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA329020	Rabbit	Human, Mouse, Rat Predicted Reactivity: Pig(100%), Bovine(100%), Horse(100%), Sheep(100%), Rabbit(100%), Dog(100%), Chicken(100%), Xenopus(100%)	ELISA

PRODUCT SYNONYMS

Frizzled 4; Polyclonal Antibody; Frizzled 4 Antibody; CD 344; CD 344 antigen; CD344; CD344 antigen; EVR1; FEVR; Frizzled (Drosophila) homolog 4; Frizzled family receptor 4; Frizzled homolog 4 (Drosophila); Frizzled-4; Fz 4; Fz-4; Fz4; FZD4; FZD4_HUMAN; FZD4S; FzE4; GPCR; hFz4; MGC34390; WNT receptor frizzled 4; anti-FZD4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat Predicted Reactivity: Pig(100%), Bovine(100%), Horse(100%), Sheep(100%), Rabbit(100%), Dog(100%), Chicken(100%), Xenopus(100%)
Clonality	Polyclonal
Isotype	IgG
Specificity	Frizzled 4 Antibody detects endogenous levels of total Frizzled 4.
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin (Thermo Fisher Scientific).
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Concentration	1mg/ml (varies by lot)
Sequence Length	537
Applicable Applications for anti-FZD4 antibody	ELISA
Immunogen	A synthesized peptide derived from human Frizzled 4, corresponding to a region within the internal amino acids.
Fragment	Fab fragment
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



IHC (Immunohistochemisry)
AAA329020 at 1/100 staining Human kidney cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4Â°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohiostchemistry)
AAA329020 at 1/100 staining Rat heart tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4Â°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



WB (Western Blot)
Western blot analysis of extracts from B16F10 cells, using Frizzled 4 Antibody. The lane on the left was treated with blocking peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-FZD4 antibody	Function: Receptor for Wnt proteins (PubMed:30135577). Most frizzled receptors are coupled to the beta-catenin (CTNNB1) canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin (CTNNB1) and activation of Wnt target genes (PubMed:30135577). Plays a critical role in retinal vascularization by acting as a receptor for Wnt proteins and norrin (NDP) (By similarity). In retina, it can be activated by Wnt protein-binding and also by Wnt-independent signaling via binding of norrin (NDP), promoting in both cases beta-catenin (CTNNB1) accumulation and stimulation of LEF/TCF-mediated transcriptional programs (By similarity). A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Post Translational Modifications: Ubiquitinated by ZNRF3, leading to its degradation by the proteasome. Subcellular Location: Cell membrane>Multi-pass membrane protein. Tissue Specificity: Almost ubiquitous (PubMed:10544037). Largely expressed in adult heart, skeletal muscle, ovary, and fetal kidney (PubMed:10544037). Moderate amounts in adult liver, kidney, pancreas, spleen, and fetal lung, and small amounts in placenta, adult lung, prostate, testis, colon, fetal brain and liver (PubMed:10544037). Subunit Structure: Interacts with MAGI3 and NDP. Component of a complex, at least composed of TSPAN12, FZD4 and norrin (NDP). Interacts (via FZ domain) with TSKU. Interacts with glypican GPC3 (PubMed:24496449). Similarity: Lys-Thr-X-X-X-Trp motif interacts with the PDZ domain of Dvl (Disheveled) family members and is involved in the activation of the Wnt/beta-catenin signaling pathway. The FZ domain is involved in binding with Wnt ligands. Belongs to the G-protein coupled receptor Fz/Smo family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	22547161
NCBI GeneID	8322
NCBI Accession #	NP_036325.2
NCBI GenBank Nucleotide #	NP_036325.2
UniProt Accession #	Q9ULV1
Molecular Weight	Observed Molecular Weight: (Observed)59kD. Predicted Molecular Weight: (Calculated)60kDa.
NCBI Official Full Name	frizzled-4
NCBI Official Synonym Full Names	frizzled class receptor 4
NCBI Official Symbol	FZD4
NCBI Official Synonym Symbols	Fz4; EVR1; FEVR; Fz-4; FzE4; GPCR; hFz4; CD344; FZD4S
NCBI Protein Information	frizzled-4
UniProt Protein Name	Frizzled-4
UniProt Gene Name	FZD4
UniProt Synonym Gene Names	Fz-4; hFz4
UniProt Entry Name	FZD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.