

Rabbit anti-Human, Mouse Cytokeratin 18 Polyclonal Antibody | anti-KRT18 antibody

Phospho-Cytokeratin 18 (Ser53) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA329337	Rabbit	Human, Mouse	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Cytokeratin 18; Polyclonal Antibody; Phospho-Cytokeratin 18 (Ser53) Antibody; Cell proliferation inducing gene 46 protein; Cell proliferation inducing protein 46; Cell proliferation-inducing gene 46 protein; CK 18; CK-18; CK18; CYK 18; CYK18; Cytokeratin endo B; Cytokeratin-18; K 18; K18; K1C18_HUMAN; KA18; Keratin 18; type I; Keratin D; keratin; type I cytoskeletal 18; Keratin-18; Krt18; anti-KRT18 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Phospho-Cytokeratin 18 (Ser53) Antibody detects endogenous levels of Cytokeratin 18 only when phosphorylated at Ser53. Tissue Specificity: Expressed in colon, placenta, liver and very weakly in exocervix. Increased expression observed in lymphHnodes of breast carcinoma.
Purity/Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-KRT18 antibody	ELISA (Peptide ELISA), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human Cytokeratin 18 around the phosphorylation site of Ser53.
Conjugation	Unconjugated
Fragment	Fab fragment
Post Translational Modifications	Phosphorylation at Ser-34 increases during mitosis. Hyperphosphorylated at Ser-53 in diseased cirrhosis liver. Phosphorylation increases by IL-6. Proteolytically cleaved by caspases during epithelial cell apoptosis. Cleavage occurs at Asp-238 by either caspase-3, caspase-6 or caspase-7. O-GlcNAcylation increases solubility, and decreases stability by inducing proteasomal degradation.
Subunit Structure	Heterotetramer of two type I and two type II keratins. KRT18 associates with KRT8. Interacts with the thrombin-antithrombin complex (By similarity). Interacts with PNN and mutated CFTR. Interacts with YWHAE, YWHAH and YWHAZ only when phosphorylated. Interacts with DNAJB6, TCHP and TRADD. Interacts with FAM83H. Interacts with EPPK1 (By similarity). (Microbial infection) Interacts with hepatitis C virus/HCV core protein.
Similarity	Belongs to the intermediate filament family.
Subcellular Location	Cytoplasm>Perinuclear region. Nucleus>Nucleolus.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)
At 1/100 staining Human gastric cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry)
At 1/100 staining Human colorectal cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)
At 1/100 staining Human pancreatic cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)
Staining Human kidney cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



WB (Western Blot)
Western blot analysis of extracts from Mouse brain, using Phospho-Cytokeratin 18 (Ser53) Antibody. The lane on the left was treated with blocking peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT18 antibody	Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557888
NCBI GeneID	3875
NCBI Accession #	NP_000215.1
NCBI GenBank Nucleotide #	NM_000224.2
UniProt Accession #	P05783; Q53G38; Q5U0N8; Q9BW26
Molecular Weight	48,058 Da
NCBI Official Full Name	keratin, type I cytoskeletal 18
NCBI Official Synonym Full Names	keratin 18
NCBI Official Symbol	KRT18
NCBI Official Synonym Symbols	K18; CYK18
NCBI Protein Information	keratin, type I cytoskeletal 18; CK-18; keratin-18; cytokeratin 18; cytokeratin-18; cell proliferation-inducing protein 46; cell proliferation-inducing gene 46 protein
UniProt Protein Name	Keratin, type I cytoskeletal 18
UniProt Gene Name	KRT18

UniProt Synonym Gene Names	CYK18; CK-18; K18
UniProt Entry Name	K1C18_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cytokeratin-18-antibody-peptide-elisa-immunohistochemistry-western-blot-elisa/329337>