

Rabbit PEA15 Polyclonal Antibody | anti-PEA15 antibody

Phospho-PEA15 (Ser104) Antibody

CATALOG # AAA329371	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Immunohistochemistry
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

PEA15; Polyclonal Antibody; Phospho-PEA15 (Ser104) Antibody; 15kDa phosphoprotein enriched in astrocytes; Astrocytic phosphoprotein PEA 15; Astrocytic phosphoprotein PEA-15; Astrocytic phosphoprotein PEA15; HMAT 1; HMAT1; Homolog of mouse MAT 1 oncogene; Homolog of mouse MAT1 oncogene; HUMMAT 1H; HUMMAT1H; MAT 1; MAT 1H; MAT1; MAT1H; PEA 15; Pea15; PEA15 protein; PEA15_HUMAN; PED; Phosphoprotein enriched in astrocytes 15; Phosphoprotein enriched in astrocytes 15kD; Phosphoprotein enriched in diabetes; anti-PEA15 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Phospho-PEA15 (Ser104) Antibody detects endogenous levels of PEA15 only when phosphorylated at Ser104. Tissue Specificity: Ubiquitously expressed. Most abundant in tissues such as heart, brain, muscle and adipose tissue which utilize glucose as an energy source. Lower expression in glucose-producing tissues. Higher levels of expression are found in tissues from individuals with type 2 diabetes than in controls.
Purity/Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-PEA15 antibody	ELISA (Peptide ELISA), IHC (Immunohistochemistry)
Immunogen	A synthesized peptide derived from human PEA15 around the phosphorylation site of Ser104.
Conjugation	Unconjugated
Fragment	Fab fragment
Post Translational Modifications	Phosphorylated by protein kinase C and calcium-calmodulin-dependent protein kinase. These phosphorylation events are modulated by neurotransmitters or hormones.
Subunit Structure	Binds RPS6KA3, MAPK3 and MAPK1. Transient interaction with PLD1 and PLD2 (By similarity). Interacts with CASP8 and FADD.
Subcellular Location	Cytoplasm. Note: Associated with microtubules.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.


IMAGES




IHC (Immunohistochemistry)
At 1/100 staining Human ovarian cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)
At 1/100 staining Mouse liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)
Staining Mouse kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)
At 1/100 staining Rat pancreatic tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-PEA15 antibody	Blocks Ras-mediated inhibition of integrin activation and modulates the ERK MAP kinase cascade. Inhibits RPS6KA3 activities by retaining it in the cytoplasm (By similarity). Inhibits both TNFRSF6- and TNFRSF1A-mediated CASP8 activity and apoptosis. Regulates glucose transport by controlling both the content of SLC2A1 glucose transporters on the plasma membrane and the insulin-dependent trafficking of SLC2A4 from the cell interior to the surface.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505705
NCBI GeneID	8682
NCBI Accession #	NP_003759.1
NCBI GenBank Nucleotide #	NM_003768.3
UniProt Accession #	Q15121; O00511
Molecular Weight	15,040 Da
NCBI Official Full Name	astrocytic phosphoprotein PEA-15

NCBI Official Synonym Full Names	phosphoprotein enriched in astrocytes 15
NCBI Official Symbol	PEA15
NCBI Official Synonym Symbols	PED; MAT1; HMAT1; MAT1H; PEA-15; HUMMAT1H
NCBI Protein Information	astrocytic phosphoprotein PEA-15; homolog of mouse MAT-1 oncogene; phosphoprotein enriched in diabetes; Phosphoprotein enriched in astrocytes, 15kD; 15 kDa phosphoprotein enriched in astrocytes
UniProt Protein Name	Astrocytic phosphoprotein PEA-15
UniProt Gene Name	PEA15
UniProt Synonym Gene Names	PED
UniProt Entry Name	PEA15_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/pea15-antibody-peptide-elisa-immunohistochemistry-elisa/329371>