

Rabbit CD22 Polyclonal Antibody | anti-CD22 antibody

Phospho-CD22 (Tyr807) Antibody

CATALOG # AAA329373	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Immunohistochemistry
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

CD22; Polyclonal Antibody; Phospho-CD22 (Tyr807) Antibody; B cell receptor CD22 precursor; B lymphocyte cell adhesion molecule; B-cell receptor CD22; B-lymphocyte cell adhesion molecule; BL CAM; BL-CAM; BLCAM; CD 22; CD22 antigen; CD22 molecule; CD22 protein; CD22_HUMAN; Lectin 2; Leu14; Lyb8; MGC130020; sialic acid binding Ig like lectin 2; Sialic acid binding immunoglobulin like lectin 2; Sialic acid-binding Ig-like lectin 2; SIGLEC 2; Siglec-2; SIGLEC2; T cell surface antigen Leu 14; T-cell surface antigen Leu-14; anti-CD22 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Phospho-CD22 (Tyr807) Antibody detects endogenous levels of CD22 only when phosphorylated at Tyr807. Tissue Specificity: B-lymphocytes.
Purity/Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-CD22 antibody	ELISA (Peptide ELISA), IHC (Immunohistochemistry)
Immunogen	A synthesized peptide derived from human CD22 around the phosphorylation site of Tyr807.
Conjugation	Unconjugated
Fragment	Fab fragment
Post Translational Modifications	Phosphorylation of Tyr-762, Tyr-807 and Tyr-822 are involved in binding to SYK, GRB2 and SYK, respectively. Phosphorylation of Tyr-842 is involved in binding to SYK, PLCG2 and PIK3R1/PIK3R2. Phosphorylated on tyrosine residues by LYN.
Subunit Structure	Predominantly monomer of isoform CD22-beta. Also found as heterodimer of isoform CD22-beta and a shorter isoform. Interacts with PTPN6/SHP-1, LYN, SYK, PIK3R1/PIK3R2 and PLCG1 upon phosphorylation. Interacts with GRB2, INPP5D and SHC1 upon phosphorylation (By similarity). May form a complex with INPP5D/SHIP, GRB2 and SHC1.
Similarity	Contains 4 copies of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases. Belongs to the immunoglobulin superfamily. SIGLEC (sialic acid binding Ig-like lectin) family.
Subcellular Location	Cell membrane>Single-pass type I membrane protein.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IHC (Immunohistochemistry)
Staining Mouse colorectal tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry)
At 1/100 staining Rat spleen tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CD22 antibody	Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	297374825
NCBI GeneID	933
NCBI Accession #	NP_001172028.1
NCBI GenBank Nucleotide #	NM_001185099.1
UniProt Accession #	P20273; O95699; O95701; O95702; O95703; Q01665; Q92872; Q92873; Q9UQA6; Q9UQA7; Q9UQA8; F5H7U3
Molecular Weight	95,348 Da
NCBI Official Full Name	B-cell receptor CD22 isoform 2
NCBI Official Synonym Full Names	CD22 molecule
NCBI Official Symbol	CD22
NCBI Official Synonym Symbols	SIGLEC2; SIGLEC-2
NCBI Protein Information	B-cell receptor CD22; BL-CAM; CD22 antigen; T-cell surface antigen Leu-14; B-lymphocyte cell adhesion molecule; sialic acid binding Ig-like lectin 2; sialic acid-binding Ig-like lectin 2
UniProt Protein Name	B-cell receptor CD22
UniProt Gene Name	CD22
UniProt Synonym Gene Names	SIGLEC2; BL-CAM; Siglec-2
UniProt Entry Name	CD22_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd22-antibody-peptide-elisa-immunohistochemistry-elisa/329373>