

## Rabbit b-Enolase Polyclonal Antibody | anti-ENO3 antibody

b-Enolase Antibody

| CATALOG # | HOST   | REACTIVITY                                                                                                                                                      | APPLICATIONS                                                                       |
|-----------|--------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|
| AAA329395 | Rabbit | Human, Mouse<br>Predicted Reactivity: Pig (92%), Zebrafish (83%), Bovine (92%), Horse (92%), Sheep (92%), Rabbit (92%), Dog (92%), Chicken (92%), Xenopus (83%) | ELISA, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot |

### PRODUCT SYNONYMS

b-Enolase; Polyclonal Antibody; b-Enolase Antibody; 2 phospho D glycerate hydrol yase; 2 phospho D glycerate hydrolyase; 2-phospho-D-glycerate hydro-lyase; Beta enolase; Beta-enolase; ENO3; ENOB\_HUMAN; Enolase 3 (beta muscle); Enolase 3; Enolase3; GSD13; MSE; Muscle specific enolase; Muscle-specific enolase; Skeletal muscle enolase; anti-ENO3 antibody

### PRODUCT SPECIFICATIONS

|                                                |                                                                                                                                                                                                                                                                                                                                    |
|------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Host                                           | Rabbit                                                                                                                                                                                                                                                                                                                             |
| Reactivity                                     | Human, Mouse<br>Predicted Reactivity: Pig (92%), Zebrafish (83%), Bovine (92%), Horse (92%), Sheep (92%), Rabbit (92%), Dog (92%), Chicken (92%), Xenopus (83%)                                                                                                                                                                    |
| Clonality                                      | Polyclonal                                                                                                                                                                                                                                                                                                                         |
| Isotype                                        | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.                                                                                                                                                                                                 |
| Specificity                                    | b-Enolase Antibody detects endogenous levels of total b-Enolase.<br>Tissue Specificity: The alpha/alpha homodimer is expressed in embryo and in most adult tissues. The alpha/beta heterodimer and the beta/beta homodimer are found in striated muscle, and the alpha/gamma heterodimer and the gamma/gamma homodimer in neurons. |
| Purity/Purification                            | The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin                                                                                                                                                                                                                                       |
| Form/Format                                    | Liquid. Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.                                                                                                                                                                                         |
| Concentration                                  | 1mg/ml (varies by lot)                                                                                                                                                                                                                                                                                                             |
| Applicable Applications for anti-ENO3 antibody | ELISA (Peptide ELISA), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)                                                                                                                                                                                                           |
| Immunogen                                      | A synthesized peptide derived from human b-Enolase (ENO-3), corresponding to a region within C-terminal amino acids.                                                                                                                                                                                                               |
| Conjugation                                    | Unconjugated                                                                                                                                                                                                                                                                                                                       |
| Fragment                                       | Fab fragment                                                                                                                                                                                                                                                                                                                       |
| Subunit Structure                              | Mammalian enolase is composed of 3 isozyme subunits, alpha, beta and gamma, which can form homodimers or heterodimers which are cell-type and development-specific. Interacts with PNKD.                                                                                                                                           |
| Similarity                                     | Belongs to the enolase family.                                                                                                                                                                                                                                                                                                     |
| Subcellular Location                           | Cytoplasm.<br>Note: Localized to the Z line. Some colocalization with CKM at M-band (By similarity).                                                                                                                                                                                                                               |
| Preparation and Storage                        | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.                                                                                                                                                                                                 |

IMAGES



Application Data

At 25 degree C. Samples were then incubated with primary Ab(At 37 degree C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody.The nuclear counter stain is DAPI(blue).



IHC (Immunohistochemistry)

At 1/100 staining Mouse stomach tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemisry)

At 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4 degree C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



WB (Western Blot)

Western blot analysis of extracts from 3T3, HepG2 and C2C12 cells using b-Enolase(ENO-3) Antibody



WB (Western Blot)

Western blot analysis of extracts from Mouse kidney, using b-Enolase(ENO-3) Antibody. The lane on the left was treated with blocking peptide.

ADDITIONAL INFORMATION

|                                                    |                                                                                                                                                            |
|----------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Related Product Information for anti-ENO3 antibody | Appears to have a function in striated muscle development and regeneration.<br>Appears to have a function in striated muscle development and regeneration. |
|----------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|

NCBI AND UNIPROT INFORMATION

|                                  |                                                                                                  |
|----------------------------------|--------------------------------------------------------------------------------------------------|
| NCBI GI #                        | 425906077                                                                                        |
| NCBI GeneID                      | 2027                                                                                             |
| UniProt Accession #              | P13929; Q96AE2; B4DUI6; B4DUM6; D3DTL2; E7ENK8                                                   |
| Molecular Weight                 | 434                                                                                              |
| NCBI Official Full Name          | Beta-enolase                                                                                     |
| NCBI Official Synonym Full Names | enolase 3 (beta, muscle)                                                                         |
| NCBI Official Symbol             | ENO3                                                                                             |
| NCBI Official Synonym Symbols    | MSE; GSD13                                                                                       |
| NCBI Protein Information         | beta-enolase; muscle-specific enolase; skeletal muscle enolase; 2-phospho-D-glycerate hydrolyase |
| UniProt Protein Name             | Beta-enolase                                                                                     |
| UniProt Gene Name                | ENO3                                                                                             |
| UniProt Synonym Gene Names       | MSE                                                                                              |
| UniProt Entry Name               | ENOB_HUMAN                                                                                       |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)

Online datasheet: <https://stage.aaabiotech.com/polyclonal/b-enolase-antibody-peptide-elisa-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/329395>