

Rabbit Flt3/CD135 Polyclonal Antibody | anti-Flt3/CD135 antibody

Phospho-Flt3/CD135 (Tyr842) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA329455	Rabbit	Human, Mouse, Rat, Monkey Predicted Reactivity: Pig (92%), Zebrafish (100%), Bovine (92%), Horse (92%), Sheep (92%), Rabbit (100%), Dog (92%)	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot

PRODUCT SYNONYMS

Flt3/CD135; Polyclonal Antibody; Phospho-Flt3/CD135 (Tyr842) Antibody; CD 135; CD135; CD135 antigen; Fetal liver kinase 2; FL cytokine receptor; Flk 2; Flk2; Flt 3; FLT-3; Flt3; FLT3_HUMAN; FMS like tyrosine kinase 3; Fms related tyrosine kinase 3; Fms-like tyrosine kinase 3; Growth factor receptor tyrosine kinase type III; Ly-72; OTTHUMP0000004234; Receptor type tyrosine protein kinase FLT3; Stem cell tyrosine kinase 1; Stk 1; STK-1; Stk1; Tyrosine protein kinase receptor FLT3; Tyrosine-protein kinase receptor FLT3; anti-Flt3/CD135 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey Predicted Reactivity: Pig (92%), Zebrafish (100%), Bovine (92%), Horse (92%), Sheep (92%), Rabbit (100%), Dog (92%)
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Phospho-Flt3/CD135 (Tyr842) Antibody detects endogenous levels of Flt3/CD135 only when phosphorylated at Tyr842. Tissue Specificity: Detected in bone marrow, in hematopoietic stem cells, in myeloid progenitor cells and in granulocyte/macrophage progenitor cells (at protein level). Detected in bone marrow, liver, thymus, spleen and lymph node, and at low levels in kidney and pancreas. Highly expressed in T-cell leukemia.
Purity/Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-Flt3/CD135 antibody	ELISA (Peptide ELISA), ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide derived from human Flt3/CD135 around the phosphorylation site of Tyr842.
Conjugation	Unconjugated
Fragment	Fab fragment
Post Translational Modifications	N-glycosylated, contains complex N-glycans with sialic acid. Autophosphorylated on several tyrosine residues in response to FLT3LG binding. FLT3LG binding also increases phosphorylation of mutant kinases that are constitutively activated. Dephosphorylated by PTPRJ/DEP-1, PTPN1, PTPN6/SHP-1, and to a lesser degree by PTPN12. Dephosphorylation is important for export from the endoplasmic reticulum and location at the cell membrane. Rapidly ubiquitinated by UBE2L6 and the E3 ubiquitin-protein ligase SIAH1 after autophosphorylation, leading to its proteasomal degradation.
Subunit Structure	Monomer in the absence of bound FLT3LG. Homodimer in the presence of bound FLT3LG. Interacts with FIZ1 following ligand activation (By similarity). Interacts with FES, FER, LYN, FGR, HCK, SRC and GRB2. Interacts with PTPRJ/DEP-1 and PTPN11/SHP2. Interacts with RNF115 and RNF126 (By similarity). (Microbial infection) Interacts with human cytomegalovirus protein UL7.

Similarity	The juxtamembrane autoregulatory region is important for normal regulation of the kinase activity and for maintaining the kinase in an inactive state in the absence of bound ligand. Upon tyrosine phosphorylation, it mediates interaction with the SH2 domains of numerous signaling partners. In-frame internal tandem duplications (ITDs) result in constitutive activation of the kinase. The activity of the mutant kinase can be stimulated further by FLT3LG binding.Belongs to the protein kinase superfamily. Tyr protein kinase family. CSF-1/PDGF receptor subfamily.
Subcellular Location	Membrane>Single-pass type I membrane protein. Endoplasmic reticulum lumen. Note: Constitutively activated mutant forms with internal tandem duplications are less efficiently transported to the cell surface and a significant proportion is retained in an immature form in the endoplasmic reticulum lumen. The activated kinase is rapidly targeted for degradation.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)

Staining HeLa cells(4h of LPS treatment) by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25 degree C. Samples were then incubated with primary Ab(AF4427 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37 degree C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody.The nuclear counter stain is DAPI(blue).



WB (Western Blot)

Western blot analysis of extracts from COS7 EGF 200ng/ml 30', using Phospho-FLT3 (Tyr842) Antibody. Lane1 was treated with phospho-blocking peptide, Lane2 was treated with non-phospho-blocking peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-Flt3/CD135 antibody	Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine FLT3LG and regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and activation of the downstream effector MTOR. Promotes activation of RAS signaling and phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1. Promotes phosphorylation of FES, FER, PTPN6/SHP, PTPN11/SHP-2, PLCG1, and STAT5A and/or STAT5B. Activation of wild-type FLT3 causes only marginal activation of STAT5A or STAT5B. Mutations that cause constitutive kinase activity promote cell proliferation and resistance to apoptosis via the activation of multiple signaling pathways.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	121114304
NCBI GeneID	2322
NCBI Accession #	NP_004110.2
NCBI GenBank Nucleotide #	NM_004119.2
UniProt Accession #	P36888; Q13414; A0AVG9; B7ZLT7; B7ZLT8; F5H0A0
Molecular Weight	112,903 Da
NCBI Official Full Name	receptor-type tyrosine-protein kinase FLT3
NCBI Official Synonym Full Names	fms-related tyrosine kinase 3
NCBI Official Symbol	FLT3
NCBI Official Synonym Symbols	FLK2; STK1; CD135; FLK-2
NCBI Protein Information	receptor-type tyrosine-protein kinase FLT3; STK-1; CD135 antigen; FL cytokine receptor; fetal liver kinase 2; fms-like tyrosine kinase 3; stem cell tyrosine kinase 1; growth factor receptor tyrosine kinase type III
UniProt Protein Name	Receptor-type tyrosine-protein kinase FLT3

UniProt Gene Name	FLT3
UniProt Synonym Gene Names	CD135; FLK2; STK1; FLK-2; FLT-3; STK-1
UniProt Entry Name	FLT3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/flt3-cd135-antibody-peptide-elisa-immunocytochemistry-immunofluorescence-western-blot-elisa/329455>