

Rabbit SRF Polyclonal Antibody | anti-SRF antibody

SRF Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA329532	Rabbit	Human, Mouse, Rat Predicted Reactivity: Pig (100%), Bovine (100%), Rabbit (89%), Dog (100%)	ELISA, Western Blot


PRODUCT SYNONYMS

SRF; Polyclonal Antibody; SRF Antibody; c fos serum response element binding factor; c fos serum response element binding transcription factor; ELK3; ERP; MCM 1; MCM1; OTTHUMP00000039820; SAP2; Serum response factor; SRF serum response factor c fos serum response element binding transcription factor; SRF_HUMAN; anti-SRF antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat Predicted Reactivity: Pig (100%), Bovine (100%), Rabbit (89%), Dog (100%)
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	SRF Antibody detects endogenous levels of total SRF.
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-SRF antibody	ELISA (Peptide ELISA), WB (Western Blot)
Immunogen	A synthesized peptide derived from human SRF, corresponding to a region within N-terminal amino acids.
Conjugation	Unconjugated
Fragment	Fab fragment
Post Translational Modifications	Phosphorylated by PRKDC.
Subunit Structure	Binds DNA as a multimer, probably a dimer. Interacts with MRTFA, forming the SRF-MRTFA nuclear complex which binds the 5'-CArG-3' consensus motif (CArG box) on DNA via SRF. Forms a nuclear ternary complex with MRTFA and SCAI. Interacts with MRTFB. Interacts with MLLT7/FOXO4, NKX3A and SSRP1. Interacts with ARID2 (By similarity). Interacts with SRFBP1 (By similarity). Interacts with FOXK1. Interacts with LPXN. Interacts with OLFM2; the interaction promotes dissociation of SRF from the transcriptional repressor HEY2, facilitates binding of SRF to target genes and promotes smooth muscle differentiation.
Subcellular Location	Nucleus.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)
Western blot analysis of SRF in lysates of MCF7 , using SRF Antibody(AF4703).



WB (Western Blot)
Western blot analysis of extracts from various samples, using SRF Antibody.Lane 1: RAW264.7 cells, blocked with antigen-specific peptides,Lane 2: RAW264.7 cells,Lane 3: MCF7 cells.

ADDITIONAL INFORMATION

Related Product Information for anti-SRF antibody	SRF is a transcription factor that binds to the serum response element (SRE), a short sequence of dyad symmetry located 300 bp to the 5' of the site of transcription initiation of some genes (such as FOS). Together with MRTFA transcription coactivator, controls expression of genes regulating the cytoskeleton during development, morphogenesis and cell migration. The SRF-MRTFA complex activity responds to Rho GTPase-induced changes in cellular globular actin (G-actin) concentration, thereby coupling cytoskeletal gene expression to cytoskeletal dynamics. Required for cardiac differentiation and maturation.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507205
NCBI GeneID	6722
NCBI Accession #	NP_003122.1
NCBI GenBank Nucleotide #	NM_003131.2
UniProt Accession #	P11831; Q5T648
Molecular Weight	51,593 Da
NCBI Official Full Name	serum response factor
NCBI Official Synonym Full Names	serum response factor (c-fos serum response element-binding transcription factor)
NCBI Official Symbol	SRF
NCBI Official Synonym Symbols	MCM1
NCBI Protein Information	serum response factor
UniProt Protein Name	Serum response factor
UniProt Gene Name	SRF
UniProt Synonym Gene Names	SRF
UniProt Entry Name	SRF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

