

Rabbit anti-Human SIRT6 Polyclonal Antibody | anti-SIRT6 antibody

SIRT6 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA329578	Rabbit	Human	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

SIRT6; Polyclonal Antibody; SIRT6 Antibody; 2810449N18Rik; AI043036; Mono ADP ribosyltransferase sirtuin 6; NAD-dependent protein deacetylase sirtuin-6; Regulatory protein SIR2 homolog 6; Regulatory protein SIR2 homolog; SIR2 like 6; SIR2 like protein 6; Sir2 related protein type 6; SIR2-like protein 6; SIR2L6; SIR6_HUMAN; SIRT 6; Sirtuin (silent mating type information regulation 2 homolog) 6 (*S. cerevisiae*); Sirtuin 6; Sirtuin type 6; Sirtuin6; anti-SIRT6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	SIRT6 Antibody detects endogenous levels of total SIRT6.
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-SIRT6 antibody	ELISA (Peptide ELISA), ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide derived from human SIRT6.
Conjugation	Unconjugated
Fragment	Fab fragment
Subunit Structure	Interacts with RELA. Interacts with RBBP8; the interaction deacetylates RBBP8.
Similarity	Belongs to the sirtuin family. Class IV subfamily.
Subcellular Location	Nucleus>Nucleoplasm. Note: Predominantly nuclear. Associated with telomeric heterochromatin regions.
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



IF (Immunofluorescence)

Staining HeLa cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. Samples were then incubated with primary Ab(AF6570) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37 degree C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody. The nuclear counter stain is DAPI (blue).



WB (Western Blot)

Western blot analysis of extracts from 293 cells(heat shock treatment), using SIRT6 Antibody. The lane on the left was treated with blocking peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-SIRT6 antibody	NAD-dependent protein deacetylase. Has deacetylase activity towards histone H3K9Ac and H3K56Ac. Modulates acetylation of histone H3 in telomeric chromatin during the S-phase of the cell cycle. Deacetylates histone H3K9Ac at NF-kappa-B target promoters and may down-regulate the expression of a subset of NF-kappa-B target genes. Acts as a corepressor of the transcription factor HIF1A to control the expression of multiple glycolytic genes to regulate glucose homeostasis. Required for genomic stability. Regulates the production of TNF protein. Has a role in the regulation of life span (By similarity). Deacetylation of nucleosomes interferes with RELA binding to target DNA. May be required for the association of WRN with telomeres during S-phase and for normal telomere maintenance. Required for genomic stability. Required for normal IGF1 serum levels and normal glucose homeostasis. Modulates cellular senescence and apoptosis. On DNA damage, promotes DNA end resection via deacetylation of RBBP8. Has very weak deacetylase activity and can bind NAD (+) in the absence of acetylated substrate.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	38258612
NCBI GeneID	51548
UniProt Accession #	Q8N6T7; O75291; Q6IAF5; Q6PK99; Q8NCD2; Q9BSI5; Q9BWP3; Q9NRC7; Q9UQD1; B2RCD0
Molecular Weight	355
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-6
NCBI Official Synonym Full Names	sirtuin 6
NCBI Official Symbol	SIRT6
NCBI Official Synonym Symbols	SIR2L6
NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-6; sirtuin type 6; SIR2-like protein 6; sir2-related protein type 6; regulatory protein SIR2 homolog 6; NAD-dependent deacetylase sirtuin-6
UniProt Protein Name	NAD-dependent protein deacetylase sirtuin-6
UniProt Gene Name	SIRT6
UniProt Synonym Gene Names	SIR2L6
UniProt Entry Name	SIR6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/sirt6-antibody-peptide-elisa-immunocytochemistry-immunofluorescence-western-blot-elisa/329578>