

Mouse Xanthine Dehydrogenase/Oxidase (XDH) ELISA Kit | XDH elisa kit

Mouse Xanthine Dehydrogenase/Oxidase (XDH) ELISA Kit

CATALOG #
AAA332020

REACTIVITY
Mouse


PRODUCT SYNONYMS

Xanthine Dehydrogenase/Oxidase (XDH); Mouse Xanthine Dehydrogenase/Oxidase (XDH) ELISA Kit; XDH; XO; XOR; xanthene dehydrogenase; xanthine oxidase; xanthine oxidoreductase; XDH elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	Mouse Xanthine dehydrogenase/oxidase (XDH) ELISA Kit has high sensitivity and excellent specificity for detection of Mouse XDH. No significant cross-reactivity or interference between Mouse XDH and analogues was observed.
Sequence Length	1335
Assay Type	Sandwich ELISA (Quantitative)
Samples	Cell Culture Supernatants, Serum, Plasma, Other Biological Fluids
Detection Method	Colorimetric
Usage Notes	* Do not mix components from different kit lots or use reagents beyond the kit expiration date. * Allow all reagents to warm to room temperature for at least 30 minutes before opening. * Pre-rinse the pipet tip with reagent, use fresh pipet tips for each sample, standard and reagent to avoid contamination. * Unused wells must be kept desiccated at 4 degree C in the sealed bag provided. * Mix Thoroughly is very important for the result. It is recommended using low frequency oscillator or slight hand shaking every 10 minutes. * It is recommended that all samples and standards be assayed in duplicate or triplicate.
Preparation and Storage	Store at 2-8 degree C.

IMAGES



Standard Curve (Sample)

Fig.1. Mouse Xanthine dehydrogenase/oxidase (XDH) Standard Curve.

ADDITIONAL INFORMATION

Related Product Information for XDH elisa kit	Description: This Mouse Xanthine dehydrogenase/oxidase (XDH) ELISA Kit employs a two-site sandwich ELISA to quantitate XDH in samples. An antibody specific for XDH has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any XDH present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for XDH is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of XDH bound in the initial step. The color development is stopped and the intensity of the color is measured. Background: Xanthine dehydrogenase belongs to the group of molybdenum-containing hydroxylases involved in the oxidative metabolism of purines. The enzyme is a homodimer. Xanthine dehydrogenase can be converted to xanthine oxidase by reversible sulfhydryl oxidation or by irreversible proteolytic modification. The three substrates of this enzyme are xanthine, NAD⁺, and H₂O, whereas its three products are urate, NADH, and H⁺. This enzyme belongs to the family of oxidoreductases, to be specific, those acting on CH or CH₂ group with NAD⁺ or NADP⁺ as acceptor. Defects in xanthine dehydrogenase cause xanthinuria, may contribute to adult respiratory stress syndrome, and may potentiate influenza infection through an oxygen metabolite-dependent mechanism.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	77682555
NCBI GeneID	22436
NCBI Accession #	NP_035853.2
NCBI GenBank Nucleotide #	NP_035853.2
UniProt Accession #	Q00519
NCBI Official Full Name	xanthine dehydrogenase/oxidase
NCBI Official Synonym Full Names	xanthine dehydrogenase
NCBI Official Symbol	Xdh
NCBI Official Synonym Symbols	XO; Xor; Xox1; Xox-1
NCBI Protein Information	xanthine dehydrogenase/oxidase
UniProt Protein Name	Xanthine dehydrogenase/oxidase
UniProt Gene Name	Xdh
UniProt Synonym Gene Names	XO; XOR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.