

Mouse Transforming Growth Factor beta2 (TGF-beta2) ELISA Kit | TGF-beta2 elisa kit

Mouse Transforming Growth Factor beta2 (TGF-beta2) ELISA Kit

CATALOG #
AAA332378

REACTIVITY
Mouse


PRODUCT SYNONYMS

Transforming Growth Factor beta2 (TGF-beta2); Mouse Transforming Growth Factor beta2 (TGF-beta2) ELISA Kit; TGFB2; MGC116892; TGF-beta2; TGF-beta2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	Mouse Transforming growth factor beta2 (TGF-beta2) ELISA Kit has high sensitivity and excellent specificity for detection of Mouse TGFB2. No significant cross-reactivity or interference between Mouse TGFB2 and analogues was observed.
Sequence Length	414
Assay Type	Sandwich ELISA (Quantitative)
Samples	Cell Culture Supernatants, Serum, Plasma, Other Biological Fluids
Detection Method	Colorimetric
Usage Notes	* Do not mix components from different kit lots or use reagents beyond the kit expiration date. * Allow all reagents to warm to room temperature for at least 30 minutes before opening. * Pre-rinse the pipet tip with reagent, use fresh pipet tips for each sample, standard and reagent to avoid contamination. * Unused wells must be kept desiccated at 4 degree C in the sealed bag provided. * Mix Thoroughly is very important for the result. It is recommended using low frequency oscillator or slight hand shaking every 10 minutes. * It is recommended that all samples and standards be assayed in duplicate or triplicate.
Preparation and Storage	Store at 2-8 degree C.

IMAGES



Standard Curve (Sample)

Fig.1. Mouse Transforming growth factor beta2 (TGF-beta2) Standard Curve.

ADDITIONAL INFORMATION

Related Product Information for TGF-beta2 elisa kit	Description: This Mouse Transforming growth factor beta2 (TGF-beta2) ELISA Kit employs a two-site sandwich ELISA to quantitate TGFB2 in samples. An antibody specific for TGFB2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any TGFB2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TGFB2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TGFB2 bound in the initial step. The color development is stopped and the intensity of the color is measured. Background: Tgfb2 encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGF-beta family members. Mice lacking a functional copy of this gene display developmental defects in multiple organs and perinatal lethality. Heterozygous mutant mice exhibit aortic root aneurysm. Tgfb2 encodes multiple isoforms that may undergo similar proteolytic processing.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	342187041
NCBI GeneID	21808
UniProt Accession #	P27090
NCBI Official Full Name	Transforming growth factor beta-2 proprotein
NCBI Official Synonym Full Names	transforming growth factor, beta 2
NCBI Official Symbol	Tgfb2
NCBI Official Synonym Symbols	Tgfb-2; BB105277; Tgf-beta2
NCBI Protein Information	transforming growth factor beta-2 proprotein; transforming growth factor beta-2
UniProt Protein Name	Transforming growth factor beta-2
UniProt Gene Name	Tgfb2
UniProt Synonym Gene Names	TGF-beta-2; LAP
UniProt Entry Name	TGFB2_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.