

Human IL-17 ELISA Kit | IL-17 elisa kit

Human IL-17 ELISA Kit

CATALOG # AAA332551	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

IL-17; Human IL-17 ELISA Kit; IL 17; IL17A; CTLA8; IL-17A; IL17; cytotoxic T-lymphocyte-associated antigen 8; cytotoxic T-lymphocyte-associated protein 8; cytotoxic T-lymphocyte-associated serine esterase 8; interleukin 17 (cytotoxic T-ly; IL-17 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	Human IL-17 ELISA Kit has high sensitivity and excellent specificity for detection of Human IL-17. No significant cross-reactivity or interference between Human IL-17 and analogues was observed.
Assay Type	Sandwich
Samples	Serum, Plasma, Cell culture supernatants
Detection Range	7.8 pg/mL-500 pg/mL
Sensitivity	4 pg/mL
Applicable Applications for IL-17 elisa kit	ELISA
Intra-assay Precision	Intra-assay Precision: The CV (%)?10%.
Inter-assay Precision	Inter-assay Precision :The CV (%)?12%
Preparation and Storage	The unopened kit should be stored at 2 - 8 degree C. After opening, please store refer to protocols.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL-17 elisa kit	Principle of the Assay: The protein encoded by IL-17 gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). Human IL-17 ELISA Kit employs a double antibody sandwich method to quantitate Human IL-17 in samples. An antibody specific for Human IL-17 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any Human IL-17 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for Human IL-17 is added to the wells. After washing, proprietary Streptavidin-HRP conjugates is added to the wells. Following a wash to remove any unbound streptavidin-enzyme reagent, adding HRP Substrate (TMB), TMB turns blue under the catalysis of HRP, and turns yellow after adding stop solution. Measure the OD value with a microplate reader at 450nm wavelength. The IL-17 concentration is proportional to the OD450 nm value.
Product Categories/Family for IL-17 elisa kit	ELISA Assay Kit; Protein

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504651
NCBI GeneID	3605
NCBI Accession #	NP_002181.1
NCBI GenBank Nucleotide #	NM_002190.2
UniProt Accession #	Q16552; Q5T2P0
Molecular Weight	17,504 Da
NCBI Official Full Name	interleukin-17A
NCBI Official Synonym Full Names	interleukin 17A
NCBI Official Symbol	IL17A
NCBI Official Synonym Symbols	IL17; CTLA8; IL-17; IL-17A
NCBI Protein Information	interleukin-17A; CTLA-8; cytotoxic T-lymphocyte-associated antigen 8; cytotoxic T-lymphocyte-associated protein 8; interleukin 17 (cytotoxic T-lymphocyte-associated serine esterase 8)
UniProt Protein Name	Interleukin-17A
UniProt Gene Name	IL17A
UniProt Synonym Gene Names	CTLA8; IL17; IL-17; IL-17A; CTLA-8
UniProt Entry Name	IL17_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/il-17/332551>