

Mouse Myc-Tag Monoclonal Antibody | anti-MYC-Tag antibody

Myc-Tag Monoclonal Antibody

CATALOG # AAA36625	HOST Mouse	APPLICATIONS Immunoprecipitation, Immunofluorescence, Western Blot
------------------------------	----------------------	--


PRODUCT SYNONYMS

Myc-Tag; Monoclonal Antibody; Myc-Tag Monoclonal Antibody; Mouse anti c-myc mAb: N/A; anti-MYC-Tag antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-MYC-Tag antibody	IP (Immunoprecipitation), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthetic peptide of Myc-Tag.
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



WB (Western Blot)

Western blot analysis of over-expressed Myc-tagged protein in 293T cell lysate, using Myc-Tag antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-MYC-Tag antibody	Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. The c-Myc tag corresponds to amino acid residues(EQKLISEEDL) of the human c-Myc protein. It can be used for affinity chromatography, then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits. Myc-Tag mouse mAb detects recombinant proteins containing the Myc tag. The antibody recognizes the Myc-tag EQKLISEEDL fused to either the amino- or carboxy- terminus of targeted proteins.
Product Categories/Family for anti-MYC-Tag antibody	Tag mAb

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/myc-tag-antibody-immunoprecipitation-immunofluorescence-western-blot-elisa/36625>