

Rabbit NCAM1 Polyclonal Antibody | anti-NCAM1 antibody

NCAM1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA36666	Rabbit	Human, Mouse, Rat	Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

NCAM1; Polyclonal Antibody; NCAM1 Polyclonal Antibody; CD56\NCAM: CD56; MSK39; NCAM; anti-NCAM1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	858
Applicable Applications for anti-NCAM1 antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human NCAM1
Calculated Molecular Weight	95kD
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



WB (Western Blot)

Western blot analysis of extracts of mouse brain, using NCAM1 antibody. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-NCAM1 antibody	Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the immunoglobulin (Ig) superfamily. It is a multifunction protein involved in synaptic plasticity, neurodevelopment, and neurogenesis. NCAM1 is expressed on human neurones, glial cells, skeletal muscle cells, NK cells and a subset of T cells, and the expression is observed in a wide variety human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronectin-type III repeats (FNIII). All three forms can be posttranslationally modified by addition of polysialic acid (PSA) (PMID: 14976519). Several other isoforms have also been described.
Product Categories/Family for anti-NCAM1 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	205830665
NCBI GeneID	4684
UniProt Accession #	P13591; P13592; P13593; Q05C58; Q15829; Q16180; Q16209; Q59FL7; Q86X47; Q96CJ3; A8K8T8
Molecular Weight	858
NCBI Official Full Name	Neural cell adhesion molecule 1
NCBI Official Synonym Full Names	neural cell adhesion molecule 1
NCBI Official Symbol	NCAM1
NCBI Official Synonym Symbols	CD56; NCAM; MSK39
NCBI Protein Information	neural cell adhesion molecule 1; neural cell adhesion molecule, NCAM; antigen recognized by monoclonal antibody 5.1H11
UniProt Protein Name	Neural cell adhesion molecule 1
UniProt Gene Name	NCAM1
UniProt Synonym Gene Names	NCAM; N-CAM-1; NCAM-1
UniProt Entry Name	NCAM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ncam1-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/36666>