

Rabbit anti-Human, Mouse CRP Polyclonal Antibody | anti-CRP antibody

CRP Polyclonal Antibody

CATALOG # AAA36681	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot
------------------------------	-----------------------	-----------------------------------	---


PRODUCT SYNONYMS

CRP; Polyclonal Antibody; CRP Polyclonal Antibody; MGC149895; MGC88244; PTX1; anti-CRP antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.
Concentration	1mg/ml (varies by lot)
Sequence Length	224
Applicable Applications for anti-CRP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human CRP
Calculated Molecular Weight	25kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IF (Immunofluorescence)
Immunofluorescence analysis of HeLa cells using CRP antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)
Western blot analysis of extracts of rat brain, using CRP antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-CRP antibody	Human c-Reactive protein (CRP) was discovered in serum of patients as a nonantibody precipitin of the C-polysaccharide from pneumococci(1). During acute infection, the amounts of circulating CPR increases 100-1000 fold (2). Serum CRP values are routinely measured to detect or monitor many human diseases and minor elevations of C-reactive protein are predictive of cardiovascular events in patients with coronary heart disease (3). CRP promotes host defense, is an efficient activator of the classical C cascade, and shown to be opsonic (4).The opsonic and C activation activities are responsible for the nonspecific host defense activities providing protection in the preimmune stage of acute infections. Additionally, IL-1 and IL-6 are known to stimulate CRP biosynthesis (5).
Product Categories/Family for anti-CRP antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	117486
NCBI GeneID	1401
UniProt Accession #	P02741; Q08AK3; Q8WW75; A8K078; D3DVD9; D3DVE0
Molecular Weight	224
NCBI Official Full Name	C-reactive protein
NCBI Official Synonym Full Names	C-reactive protein, pentraxin-related
NCBI Official Symbol	CRP
NCBI Official Synonym Symbols	PTX1
NCBI Protein Information	C-reactive protein; pentraxin 1
UniProt Protein Name	C-reactive protein
UniProt Gene Name	CRP
UniProt Synonym Gene Names	PTX1
UniProt Entry Name	CRP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

