

Rabbit NSF Polyclonal Antibody | anti-NSF antibody

Rabbit polyclonal to Human NSF

CATALOG # AAA36735	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

NSF; Polyclonal Antibody; Rabbit polyclonal to Human NSF; NSF Rabbit pab Antibody; anti-NSF antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purified
Form/Format	PBS with 0.09% Sodium azide, 50% glycerol, pH 7.3.
Concentration	1mg/ml (varies by lot)
Sequence Length	744
Applicable Applications for anti-NSF antibody	WB (Western Blot)
Immunogen	A recombinant protein of human NSF
Preparation and Storage	Upon delivery aliquot and store at -20 degree C or -80 degree C

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat heart using NSF antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver injury using NSF antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using NSF antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 15s.

ADDITIONAL INFORMATION

Related Product Information for anti-NSF antibody	Several protein-protein interactions are essential to membrane fusion during endocytosis. Membrane fusion requires interaction among SNARE1 proteins associated with both donor and acceptor membranes (1,2). Following membrane fusion, the alpha-SNAP cytoplasmic adapter protein binds to the SNARE complex. N-ethylmaleimide-sensitive factor (NSF), a hexameric ATPase, then associates with the alpha-SNAP/SNARE complex to mediate SNARE disassembly during membrane fusion (3,4). The ATPase activity of NSF induces a conformational change in the alpha-SNAP/SNARE complex that leads to its dissociation from the membrane, membrane fusion and eventual recycling of the SNARE complex for subsequent membrane fusion (3,4).
References	1. Ungermann, C. and Langosch, D. (2005) J Cell Sci 118, 3819-28. 2. Leabu, M. (2006) J Cell Mol Med 10, 423-7. 3. May, A.P. et al. (2001) J Biol Chem 276, 21991-4. 4. Dalal, S. et al. (2004) Mol Biol Cell 15, 637-48.

NCBI AND UNIPROT INFORMATION

NCBI GI #	156564401
NCBI GeneID	4905
NCBI Accession #	NP_006169.2
NCBI GenBank Nucleotide #	NM_006178.3
UniProt Accession #	P46459 ; Q8N6D7 ; Q9UKZ2 ; A8K2D9
Molecular Weight	82,594 Da
NCBI Official Full Name	vesicle-fusing ATPase
NCBI Official Synonym Full Names	N-ethylmaleimide-sensitive factor
NCBI Official Symbol	NSF
NCBI Official Synonym Symbols	SKD2
NCBI Protein Information	vesicle-fusing ATPase; NEM-sensitive fusion protein; vesicular-fusion protein NSF; N-ethylmaleimide-sensitive fusion protein; N-ethylmaleimide-sensitive factor-like protein
UniProt Protein Name	Vesicle-fusing ATPase
UniProt Gene Name	NSF
UniProt Synonym Gene Names	NEM-sensitive fusion protein

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nsf-antibody-western-blot-elisa/36735>