

Rabbit Phospho-Stat1(pS727) Antibody | anti-STAT1 antibody

Phospho-Stat1(pS727)

CATALOG # AAA36800	HOST Rabbit	APPLICATIONS Immunoprecipitation, Immunohistochemistry, Western Blot
------------------------------	-----------------------	--


PRODUCT SYNONYMS

Phospho-Stat1(pS727); STAT1; anti-STAT1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Sequence Length	750
Applicable Applications for anti-STAT1 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Cross Reactivity	Human, Mouse, Rat
Immunogen	A synthetic phosphorylated peptide around S727 of human STAT1 (NP_009330.1).
Preparation and Storage	Store at -20°C. Avoid freeze / thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human esophageal using Phospho-STAT1-S727 Rabbit pAb (AAA36800) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of HeLa, using Phospho-STAT1-S727 antibody (AP0109) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.

ADDITIONAL INFORMATION

Related Product Information for anti-STAT1 antibody	Background: The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. The protein encoded by this gene can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. The protein plays an important role in immune responses to viral, fungal and mycobacterial pathogens. Mutations in this gene are associated with Immunodeficiency 31B, 31A, and 31C.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	6274552
NCBI GeneID	6772
NCBI Accession #	NP_009330.1
NCBI GenBank Nucleotide #	NM_007315.3
UniProt Accession #	P42224; Q53S88; Q53XW4; Q68D00; Q9UDL5; A8K989; B2RCA0; D2KFR8; D3DPI7
Molecular Weight	Observed MW: 91KDa Calculated MW: 87kDa
NCBI Official Full Name	signal transducer and activator of transcription 1-alpha/beta isoform alpha
NCBI Official Synonym Full Names	signal transducer and activator of transcription 1, 91kDa
NCBI Official Symbol	STAT1
NCBI Official Synonym Symbols	CANDF7; ISGF-3; STAT91
NCBI Protein Information	signal transducer and activator of transcription 1-alpha/beta; transcription factor ISGF-3 components p91/p84; signal transducer and activator of transcription-1
UniProt Protein Name	Signal transducer and activator of transcription 1-alpha/beta
UniProt Gene Name	STAT1
UniProt Entry Name	STAT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.