

Rabbit anti-Human CTLA4 Polyclonal Antibody | anti-CTLA4 antibody

CTLA4 Polyclonal Antibody

CATALOG # AAA36834	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot
------------------------------	-----------------------	----------------------------	-------------------------------------


PRODUCT SYNONYMS

CTLA4; Polyclonal Antibody; CTLA4 Polyclonal Antibody; CD; CD152; CELIAC3; CTLA-4; GRD4; GSE; ICOS; IDDM12; anti-CTLA4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	223
Applicable Applications for anti-CTLA4 antibody	WB (Western Blot)
Species	Human
Route	Synthetic PeptideCategory
Immunogen	A synthetic peptide of human AKT1
Calculated Molecular Weight	25kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CTLA4 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 5s.

ADDITIONAL INFORMATION

Related Product Information for anti-CTLA4 antibody	This gene is a member of the immunoglobulin superfamily and encodes a protein which transmits an inhibitory signal to T cells. The protein contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. The membrane-bound isoform functions as a homodimer interconnected by a disulfide bond, while the soluble isoform functions as a monomer. Mutations in this gene have been associated with insulin-dependent diabetes mellitus, Graves disease, Hashimoto thyroiditis, celiac disease, systemic lupus erythematosus, thyroid-associated orbitopathy, and other autoimmune diseases. [provided by RefSeq, Jul 2008]
Product Categories/Family for anti-CTLA4 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	27735177
NCBI GeneID	1493
UniProt Accession #	P16410; O95653; Q0PP65; Q52MC1; Q53TD5; Q5S005; Q8WXJ1; Q96P43; Q9UKN9; A0N1S0; E9PDH0
Molecular Weight	223
NCBI Official Full Name	Cytotoxic T-lymphocyte protein 4
NCBI Official Synonym Full Names	cytotoxic T-lymphocyte-associated protein 4
NCBI Official Symbol	CTLA4
NCBI Official Synonym Symbols	CD; GSE; GRD4; ALPS5; CD152; CTLA-4; IDDM12; CELIAC3
NCBI Protein Information	cytotoxic T-lymphocyte protein 4; CD152 isoform; celiac disease 3; insulin-dependent diabetes mellitus 12; cytotoxic T-lymphocyte-associated serine esterase-4; cytotoxic T lymphocyte associated antigen 4 short spliced form; ligand and transmembrane spliced cytotoxic T lymphocyte associated antigen 4
UniProt Protein Name	Cytotoxic T-lymphocyte protein 4
UniProt Gene Name	CTLA4
UniProt Synonym Gene Names	CD152; CTLA-4
UniProt Entry Name	CTLA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.