

Rabbit anti-Human, Mouse CD55 Polyclonal Antibody | anti-CD55 antibody

CD55 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA36963	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD55; Polyclonal Antibody; CD55 Polyclonal Antibody; CR; TC; DAF; CROM; anti-CD55 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	381
Applicable Applications for anti-CD55 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human CD55
Calculated Molecular Weight	41kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of A549 cells using CD55 antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CD55 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-CD55 antibody	This gene encodes a protein involved in the regulation of the complement cascade. The encoded glycoprotein is also known as the decay-accelerating factor (DAF); binding of DAF to complement proteins accelerates their decay, disrupting the cascade and preventing damage to host cells. Antigens present on the DAF glycoprotein constitute the Cromer blood group system (CROM). Two alternatively spliced transcripts encoding different proteins have been identified. The predominant transcript encodes a membrane-bound protein expressed on cells exposed to plasma component proteins but an alternatively spliced transcript produces a soluble protein present at much lower levels. Additional, alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq, Jul 2008]
Product Categories/Family for anti-CD55 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	60416353
NCBI GeneID	1604
UniProt Accession #	P08174; P09679; P78361; Q14UF2; Q14UF3; Q14UF4; Q14UF5; Q14UF6; B1AP14; D3DT83; D3DT84; E7ER69
Molecular Weight	381
NCBI Official Full Name	Complement decay-accelerating factor
NCBI Official Synonym Full Names	CD55 molecule, decay accelerating factor for complement (Cromer blood group)
NCBI Official Symbol	CD55
NCBI Official Synonym Symbols	CR; TC; DAF; CROM
NCBI Protein Information	complement decay-accelerating factor; CD55 antigen
UniProt Protein Name	Complement decay-accelerating factor
UniProt Gene Name	CD55
UniProt Synonym Gene Names	CR; DAF
UniProt Entry Name	DAF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.